

# WAVELETS CENTERED ON A KNOT SEQUENCE: PIECEWISE POLYNOMIAL WAVELETS ON A QUASI-CRYSTAL LATTICE

BRUCE W. ATKINSON, DEREK O. BRUFF,  
JEFFREY S. GERONIMO, AND DOUGLAS P. HARDIN

**ABSTRACT.** We develop a general notion of orthogonal wavelets ‘centered’ on an irregular knot sequence. We present two families of orthogonal wavelets that are continuous and piecewise polynomial. As an application, we construct continuous, piecewise quadratic, orthogonal wavelet bases on the quasi-crystal lattice consisting of the  $\tau$ -integers where  $\tau$  is the golden-mean. The resulting spaces then generate a multiresolution analysis of  $L^2(\mathbf{R})$  with scaling factor  $\tau$ .

## INTRODUCTION

Wavelets are a useful tool for representing general functions and datasets and are now used in a wide variety of areas such as signal processing, image compression, function approximation, and finite-element methods. Traditionally wavelets are constructed from one function, the so-called mother wavelet, by integer translation and integer dilation and gives rise to a stationary refinement equation (See Daubechies [6]). For many functions however this may not be the most efficient way of approximating them and one may require adaptive schemes for which the knot sequence is not regularly spaced or where the knots are on a regular grid but the refinement equation changes at each step. There has been much work devoted to extending wavelet construction to irregularly spaced knot sequences including the lifting scheme of Sweldens [21] and wavelets on irregular grids (see Daubechies et. al. [7],[8], Charina and Stöckler [3]) or multiresolution analyses with irrational scaling factors (see Chui and Shi [4] and Hernández, Wang, and Weiss [17], [18]) or to nonstationary refinement masks (see Cohen and Dyn [5] and Herley, Kovačević, and Vetterli [16]). If the knots are allowed to be chosen generally enough it is difficult to maintain the orthogonality of the wavelet functions while keeping the compact support and smoothness properties.

In [9, 12] piecewise polynomial, orthogonal, multiwavelets with compact support were constructed using multiresolution analyses that were *intertwined* with classical spline spaces. Both the multiwavelets as well as the scaling functions generating these multiwavelets have ‘short’ support allowing them to be adapted to irregular knot sequences using the machinery of *squeezable bases* developed in [10, 11] in a way that preserved the orthogonality, polynomial reproduction, and smoothness of the respective bases. In general, the multiresolution analyses were restricted to *semi-regular* refinement schemes in which an initial irregular knot sequence is refined by a regular refinement scheme such as mid-point subdivision (cf. [2]). However, in

---

*Key words and phrases.* orthogonal wavelets; piecewise polynomial; irregular knot sequences; quasi-crystal lattice.

The research of JSG was partially supported by NSF grant DMS-0500641 and the research of DPH was partially supported by NSF grants DMS-0808093 and DMS-0934630.

[11], a construction was given of a fully irregular multiresolution analysis consisting of continuous, piecewise quadratic functions on an *arbitrary* sequence of nested knot sequences such that the multiresolution spaces have compactly supported orthogonal bases with ‘short’ support (we review and generalize this construction in Section 1.4). The resulting spaces in the multiresolution analysis did not fit, for general knot sequences, into the framework of squeezable bases. In [1], a more general notion of *bases centered on a knot sequence* was introduced that includes the squeezable bases as well as the irregular construction. In particular, necessary and sufficient conditions were given for when a space  $V$  has an *orthogonal* basis centered on a knot sequence. This is used to prove one of the main results of this paper that if  $V^0 \subset V^1$  are spaces generated by orthogonal bases centered on a common knot sequence  $\mathbf{a}$  then the orthogonal complement  $W = V^1 \ominus V^0$  is also generated by an orthogonal basis  $\Psi$  centered on  $\mathbf{a}$ .

In Section 1 we review and further elucidate the theory of bases centered on a knot sequence. In particular, as mentioned above, we focus on the question of characterizing spaces that are generated by orthogonal bases centered on a knot sequence. We next give two different constructions of nested spaces of this type. The first uses a fixed knot sequence  $\mathbf{a}$  and constructs  $V^0 \subset V^1 \subset \dots$  which are spaces generated by orthogonal continuous piecewise polynomial functions with compact support and with breakpoints in  $\mathbf{a}$ ; these spaces have increasing polynomial reproduction. The second construction begins with a given knot sequence  $\mathbf{a}$  and the spline space  $S_2^0(\mathbf{a})$  consisting of continuous piecewise quadratic functions with breakpoints in  $\mathbf{a}$ . The knot sequence  $\mathbf{a}$  is then refined and a space  $V$ , containing  $S_2^0(\mathbf{a})$ , is constructed which is generated by continuous orthogonal piecewise quadratic functions with compact support and breakpoints in the refined sequence. If  $\mathbf{a}^0 \subset \mathbf{a}^1$  and  $V^0$  (resp.  $V^1$ ) is constructed in this way starting with  $\mathbf{a}^0$  (resp.  $\mathbf{a}^1$ ), we describe general conditions under which  $V^0 \subset V^1$ .

In Section 2 it is shown how to build wavelets from the scaling functions constructed in the previous section. Certain spaces are introduced which shed light on techniques of [12]. The wavelet construction methods given in [12] are sufficiently general so that they can be used to give a decomposition of the wavelet spaces, thus providing a convenient algorithm for calculating these bases.

In Section 3 a construction of multiwavelets is carried out for the knot sequence consisting of the  $\tau$ -integers, where  $\tau = \frac{1}{2}(1 + \sqrt{5})$  is the golden mean. These multiwavelets have a scaling factor  $\tau$  and will be called  $\tau$ -multiwavelets. The lattice generated by  $\tau$  and other Pisot numbers appears in the study of quasi-crystals and powers of these numbers appear in the diffraction patterns of actual experiments.  $\tau$ -Haar wavelets were constructed in [14, 15].  $\tau$ -Haar wavelets are orthogonal and compactly supported but they are not continuous. The above construction is used to give examples of piecewise quadratic *continuous* compactly supported  $\tau$ -multiwavelets.

## 1. BASES CENTERED ON KNOT SEQUENCES

Let  $J$  be an interval in  $\mathbf{R}$  and let  $\mathbf{a} \subset J$  have no cluster point in  $J$  and such that  $\inf \mathbf{a} = \inf J$  and  $\sup \mathbf{a} = \sup J$ ; to avoid trivial cases we assume that  $\mathbf{a}$  consists of at least three elements. We refer to such a set  $\mathbf{a}$  as a **knot sequence** in  $J$  since it can be represented as the range of a strictly increasing sequence indexed by an interval  $\mathcal{I} \subset \mathbf{Z}$ .

Let  $\mathbf{a}$  be a knot sequence in  $J$ . If  $a \in \mathbf{a}$ , we define  $a_+ := \inf\{b \in \mathbf{a} | a < b\}$ ; in this case we shall refer to  $a_+$  as the **successor** of  $a$  in  $\mathbf{a}$ . We will write  $a_{++}$  for  $(a_+)_+$ . Similarly, if  $a > \inf J$ , then we define  $a_- := \sup\{b \in \mathbf{a} | b < a\}$ ; i.e.  $a_-$  is the **predecessor** of  $a$  in  $\mathbf{a}$ . We remark that  $a_+ = \infty$  if  $a = \sup J$  and  $a_- = -\infty$  if  $a = \inf J$ .

For an arbitrary collection of functions  $G \subset L^2(J)$  and an interval  $I \subset J$ , let

$$G_I := \{g \in G \mid \text{supp } \phi \subset I\}.$$

If  $\Phi \subset L^2(J)$  is a locally finite collection of functions (i.e., on any compact interval  $K \subset J$  all but a finite number of  $\phi \in \Phi$  vanish on  $K$ ) then for  $a \in \mathbf{a}$  we define

$$(1) \quad \begin{aligned} \check{\Phi}_a &:= \Phi_{[a, a_+]}, \\ \Phi_a &:= \Phi_{[a_-, a_+]} \setminus \check{\Phi}_{a_-}, \text{ and} \\ \bar{\Phi}_a &:= \Phi_a \setminus \check{\Phi}_a, \end{aligned}$$

where we define  $\check{\Phi}_{-\infty} := \emptyset$  and  $\bar{\Phi}_{\infty} := \emptyset$ . We remark that if  $J$  contains its supremum and  $a = \sup J$  then  $\Phi_a = \emptyset$  while if  $J$  contains its infimum and  $a = \inf J$  then  $\bar{\Phi}_a = \emptyset$ . The sets  $\check{\Phi}_a$ ,  $\bar{\Phi}_a$ , and  $\Phi_a$  depend on the knot sequence  $\mathbf{a}$  since  $a_+$  and  $a_-$  are defined relative to  $\mathbf{a}$ . When there is a chance of ambiguity we write  $\check{\Phi}_{a, \mathbf{a}}$ ,  $\bar{\Phi}_{a, \mathbf{a}}$ , and  $\Phi_{a, \mathbf{a}}$  to denote the knot sequence  $\mathbf{a}$  that is referenced.

We say that  $\Phi \subset L^2(J)$  is a **basis centered on the knot sequence  $\mathbf{a}$**  provided

$$(2) \quad \begin{aligned} (a) & \text{ } \Phi \text{ is locally finite,} \\ (b) & \text{ } \Phi = \bigcup_{a \in \mathbf{a}} \Phi_a, \text{ and} \\ (c) & \text{ } (\Phi_a \cup \bar{\Phi}_{a_+})|_{[a, a_+]} \text{ is a linearly independent set for all } a \in \mathbf{a}. \end{aligned}$$

For any  $\Phi$  satisfying conditions (a) and (b) of (2), let

$$S(\Phi) := \text{clos}_{L^2(J)} \text{span } (\Phi).$$

The notion of a basis centered on a knot sequence was introduced in [1] and is a generalization of bases obtained from **minimally supported generators** as defined in [9, 11].

If  $\Phi$  is a basis centered on  $\mathbf{a}$  then it follows from properties (2) that any  $f \in S(\Phi)$  has a unique representation of the form

$$(3) \quad f = \sum_{a \in \mathbf{a}} c_a \Phi_a = \sum_{a \in \mathbf{a}} (\check{c}_a \check{\Phi}_a + \bar{c}_a \bar{\Phi}_a),$$

where the convergence of the sums is in  $L^2_{\text{loc}}(J)$ . Note that  $c_a$ ,  $\check{c}_a$ , and  $\bar{c}_a$  are treated as row vectors and  $\Phi_a$ ,  $\check{\Phi}_a$ , and  $\bar{\Phi}_a$  are treated as column vectors so that  $c_a \Phi_a$  denotes the linear combination of elements  $\Phi_a$  with coefficient vector  $c_a$ ; the expressions  $\check{c}_a \check{\Phi}_a$  and  $\bar{c}_a \bar{\Phi}_a$  are interpreted similarly. It then follows from the local linear independence condition (c) that

$$(4) \quad f \in S(\Phi) \text{ and } f = 0 \text{ on } [a, a_+] \implies \bar{c}_a, \bar{c}_{a_+} \text{ and } \check{c}_a \text{ are all 0,}$$

and then (4) implies

$$(5) \quad S(\Phi)_{[a, b]} = \text{span } (\Phi_{[a, b]}) = \text{span } (\check{\Phi}_a \cup (\bigcup_{\substack{c \in \mathbf{a} \\ a < c < b}} \Phi_c)),$$

for  $a, b \in \mathbf{a}$  with  $a < b$ . Note that if there are no knots  $c$  between  $a$  and  $b$ , then the sum (resp. union) in equation (5) is zero (resp. empty).

For a linear subspace  $V \subset L^2(J)$ , let

$$V_a := V_{[a_-, a_+]} \text{ and } \check{V}_a := V_{[a, a_+]} \quad a \in \mathbf{a}.$$

Note that if  $\Phi$  is a basis centered on the knot sequence  $\mathbf{a}$  and  $V = S(\Phi)$ , then equation (5) implies that for  $a \in \mathbf{a}$ ,

$$\check{V}_a = \text{span } \check{\Phi}_a$$

and

$$V_a = \text{span } (\check{\Phi}_{a_-} \cup \check{\Phi}_a).$$

In particular it follows that each  $V_a$  is finite dimensional. We next characterize when a linear subspace  $V \subset L^2(J)$  equals  $S(\Phi)$  for some basis  $\Phi$  centered on  $\mathbf{a}$ .

**Lemma 1.** *Let  $\mathbf{a}$  be a knot sequence in an interval  $J$  and suppose  $V$  is a linear subspace of  $L^2(J)$ . Then  $V = S(\Phi)$  for some basis  $\Phi$  centered on  $\mathbf{a}$  if and only if the following three properties hold.*

- ( $\hat{a}$ )  $V_a$  is finite dimensional for  $a \in \mathbf{a}$ .
- ( $\hat{b}$ )  $V = \text{clos}_{L^2(J)} \text{span } \left( \bigcup_{a \in \mathbf{a}} V_a \right)$ .
- ( $\hat{c}$ ) If  $f \in V$  vanishes on  $[a, a_+]$ , then  $f \in V_{(-\infty, a]} + V_{[a_+, \infty)}$ .

If ( $\hat{a}$ - $\hat{c}$ ) hold, then, for  $a \in \mathbf{a}$ , let  $\check{\Phi}_a$  be a basis for  $\check{V}_a$  and let  $\bar{\Phi}_a$  augment  $\check{\Phi}_{a_-} \cup \check{\Phi}_a$  to a basis of  $V_a$ . Then  $\Phi := \bigcup_{a \in \mathbf{a}} (\bar{\Phi}_a \cup \check{\Phi}_a)$  is a basis centered on  $\mathbf{a}$  such that  $V = S(\Phi)$ . Furthermore, any  $\Phi$  such that  $V = S(\Phi)$  is of this form, that is,  $\check{\Phi}_a$  must be a basis of  $\check{V}_a$  and  $\bar{\Phi}_a \cup \check{\Phi}_{a_-} \cup \check{\Phi}_a$  must be a basis of  $V_a$ .

*Remark 1.* Lemma 1 implies that if  $\Phi$  is a basis centered on a knot sequence  $\mathbf{b}$  that is a **refinement** of  $\mathbf{a}$  (i.e.,  $\mathbf{a} \subset \mathbf{b}$ ), then  $\Phi$  is also a basis centered on  $\mathbf{a}$ . Note that, for  $a \in \mathbf{a} \subset \mathbf{b}$ , the sets  $\Phi_{a, \mathbf{a}}$  and  $\Phi_{a, \mathbf{b}}$  are, in general, not equal.

*Proof.* ( $\Rightarrow$ ) Suppose  $V = S(\Phi)$  for some basis  $\Phi$  centered on the knot sequence  $\mathbf{a}$ . Conditions ( $\hat{a}$ ) and ( $\hat{b}$ ) then follow from (5) and condition ( $\hat{c}$ ) follows from (4).

( $\Leftarrow$ ) Suppose that ( $\hat{a}$ - $\hat{c}$ ) hold and that  $\check{\Phi}_a$  and  $\bar{\Phi}_a$  are constructed as in the statement of Lemma 1. Then ( $\hat{b}$ ) implies  $V = S(\Phi)$  and so it remains to show the local linear independence condition of  $\Phi$ . Let  $a \in \mathbf{a}$  and suppose that  $\bar{c}_a$ ,  $\check{c}_a$ , and  $\check{c}_{a_+}$  are vectors so that

$$(6) \quad f := \bar{c}_a \bar{\Phi}_a + \check{c}_a \check{\Phi}_a + \check{c}_{a_+} \bar{\Phi}_{a_+} = 0 \text{ on } [a, a_+].$$

It now suffices to show that these coefficient vectors must all vanish. By condition ( $\hat{c}$ ), we have  $f = f_1 + f_2$  where  $f_1 \in V_{(-\infty, a]}$  and  $f_2 \in V_{[a_+, \infty)}$ . Since  $f$  and  $f_2$  vanish on  $(-\infty, a_-]$  it follows that  $f_1 \in V_{[a_-, a]}$ . Hence,  $f_1 = \check{d}_{a_-} \check{\Phi}_{a_-}$  for some vector  $\check{d}_{a_-}$  by the construction of  $\check{\Phi}_{a_-}$ . Then  $g := \bar{c}_a \bar{\Phi}_a - \check{d}_{a_-} \check{\Phi}_{a_-} = 0$  on  $[a_-, a]$ . Since the support of  $g$  is in  $[a, a_+]$ , then  $g = \check{d}_a \check{\Phi}_a$  for some vector  $\check{d}_a$ , i.e.,

$$\bar{c}_a \bar{\Phi}_a - \check{d}_{a_-} \check{\Phi}_{a_-} - \check{d}_a \check{\Phi}_a = 0.$$

Since  $\bar{\Phi}_{a_-}$ ,  $\check{\Phi}_{a_-}$  and  $\check{\Phi}_a$  are linearly independent, it follows that  $\bar{c}_a = 0$ . Similarly, we have  $\bar{c}_{a_+} = 0$  and thus, by the linear independence of  $\check{\Phi}_a$  we have  $\check{c}_a = 0$ .  $\square$

**1.1. Example.** Let  $\Pi^d$  denote the space of univariate polynomials of degree at most  $d$ . For a knot sequence  $\mathbf{a}$  on an interval  $J$  and integers  $r < d$ , let

$$S_d^r(\mathbf{a}) := \{f \mid \text{supp } f \subset J, f \in C^r(J) \cap L^2(J), \text{ and } f|_{(a,a_+)} \in \Pi^d \text{ for } a \in \mathbf{a}\}$$

denote the classical spline space of degree  $d$  and regularity  $r$  intersected with  $L^2(J)$ . If  $d \geq 2r + 1$ , then  $S_d^r(\mathbf{a}) = S(\Phi)$  for a basis  $\Phi = \Phi_{r,d}^{\mathbf{a}}$  centered on  $\mathbf{a}$  (for example, the B-spline basis for  $S_d^r(\mathbf{a})$  is a basis centered on  $\mathbf{a}$ , cf. [20]). Furthermore,  $|\check{\Phi}_a| = d - 2r - 1$  if  $a < \sup J$ , and for  $\inf J < a < \sup J$ , we have  $|\Phi_a| = d - r$  and  $|\bar{\Phi}_a| = r + 1$ .

**1.2. Orthogonality Condition.** Bases  $\Phi$  and  $\Psi$  centered on the same knot sequence  $\mathbf{a}$  in the interval  $J$  are called **equivalent** if  $S(\Phi) = S(\Psi)$ . We next give a necessary and sufficient condition for a basis  $\Phi$  centered on  $\mathbf{a}$  to be equivalent to some *orthogonal* basis  $\Omega$  centered on  $\mathbf{a}$ . This condition is the main tool we use to construct orthogonal bases centered on a knot sequence. Previous versions of this lemma appeared in [9] (for the shift-invariant setting), in [11] (for the setting of “squeezeable” orthogonal bases) and in [1] (for the current setting of bases centered on a knot sequence).

**Lemma 2.** *Let  $J$  be an interval in  $\mathbf{R}$ ,  $\mathbf{a}$  a knot sequence on  $J$ ,  $\Phi$  a basis centered on  $\mathbf{a}$ , and  $V = S(\Phi)$ . Then there exists an orthogonal basis  $\Omega$  centered on the knot sequence  $\mathbf{a}$  such that  $S(\Omega) = S(\Phi)$  if and only if*

$$(7) \quad \left( I - P_{\check{V}_a} \right) V_a \perp V_{a_+} \text{ for } a \in \mathbf{a}.$$

If (7) holds, then, for  $a \in \mathbf{a}$ , let  $\check{\Omega}_a$  be an orthogonal basis of  $\check{V}_a$ ,  $\bar{\Omega}_a$  be an orthogonal basis of  $\left( I - P_{\check{V}_{a_-} \oplus \check{V}_a} \right) V_a$ , and  $\Omega_a = \check{\Omega}_a \cup \bar{\Omega}_a$ . Then  $\Omega = \{\Omega_a\}_{a \in \mathbf{a}}$  is such an orthogonal basis.

*Proof.* Suppose  $a \in \mathbf{a}$ . Note that  $\check{V}_{a_-} \perp V_{a_+}$  since their supports intersect in at most one point. It then follows that (7) is equivalent to

$$\left( I - P_{\check{V}_{a_-} \oplus \check{V}_a} \right) V_a \perp V_{a_+}.$$

Furthermore, since  $\left( I - P_{\check{V}_{a_-} \oplus \check{V}_a} \right) V_a$  is orthogonal to  $\check{V}_a$  and, due to support properties, is also orthogonal to  $\check{V}_{a_+}$ , it follows that (7) is equivalent to

$$(8) \quad \left( I - P_{\check{V}_{a_-} \oplus \check{V}_a} \right) V_a \perp \left( I - P_{\check{V}_a \oplus \check{V}_{a_+}} \right) V_{a_+}.$$

( $\Rightarrow$ ) Suppose that  $V = S(\Omega)$ , where  $\Omega$  is an orthogonal basis centered on the knot sequence  $\mathbf{a}$ . It follows from (5) and the orthogonality of  $\Omega$  that

$$V_a = \text{span } \check{\Omega}_{a_-} \oplus \text{span } \bar{\Omega}_a \oplus \text{span } \check{\Omega}_a = \check{V}_{a_-} \oplus \text{span } \bar{\Omega}_a \oplus \check{V}_a,$$

and so  $\text{span}(\bar{\Omega}_a) = \left( I - P_{\check{V}_{a_-} \oplus \check{V}_a} \right) V_a$ . Since  $\Omega$  is an orthogonal basis,  $\text{span}(\bar{\Omega}_a) \perp \text{span}(\bar{\Omega}_{a_+})$ , and so (8), and hence (7), holds.

( $\Leftarrow$ ) Suppose that  $V$  satisfies (7), and hence satisfies (8). Construct  $\Omega$  as in the statement of Lemma 2. By Lemma 1, it follows that  $\Omega$  is a basis centered on  $\mathbf{a}$  such that  $V = S(\Omega)$ . Equation (8) implies that  $\bar{\Omega}_a \perp \bar{\Omega}_{a_+}$  for  $a \in \mathbf{a}$ , which shows  $\Omega$  is an orthogonal basis of  $V$ .  $\square$

Suppose  $V = S(\Phi)$  for some  $\Phi$ . Then it is easy to verify that (7) holds if and only if

$$(9) \quad (I - P_{\check{V}_a}) \bar{\Phi}_a \perp \bar{\Phi}_{a_+}, \quad a \in \mathbf{a}.$$

We use this equation as the basis for a construction of orthogonal bases centered on a knot sequence as we next describe. Suppose  $Z = \sum_{a \in \mathbf{a}} Z_a$  where  $Z_a$ ,  $a \in \mathbf{a}$  is a finite dimensional subspace of  $L^2(J)$  such that (a) the elements of  $Z_a$  are supported in  $[a, a_+]$  and (b)  $Z_a$  is linearly independent of  $V$  restricted to  $[a, a_+]$ . Then  $U := V + Z$  satisfies the hypotheses of Lemma 1 and, by this lemma,  $U = S(\Theta)$  where  $\Theta$  is a basis centered on  $\mathbf{a}$  such that  $\check{\Theta}_a$  is a basis for  $\check{U}_a = \check{V}_a + Z_a$  and  $\bar{\Theta}_a := \bar{\Phi}_a$  for  $a \in \mathbf{a}$ . By Lemma 2,  $U = S(\Omega)$  for some orthogonal basis  $\Omega$  centered on  $\mathbf{a}$  if and only if

$$(10) \quad (I - P_{\check{V}_a + Z_a}) \bar{\Phi}_a \perp \bar{\Phi}_{a_+}, \quad a \in \mathbf{a}.$$

Without loss of generality, we may choose  $Z_a$  orthogonal to  $\check{V}_a$ . Then (10) is equivalent to

$$(11) \quad \langle (I - P_{\check{V}_a}) \bar{\Phi}_a, \bar{\Phi}_{a_+} \rangle = \langle P_{Z_a} \bar{\Phi}_a, \bar{\Phi}_{a_+} \rangle, \quad a \in \mathbf{a}.$$

where for finite collections  $F, G \subset L^2(\mathbf{R})$ , we let  $\langle F, G \rangle$  denote the matrix  $(\langle f, g \rangle)_{f \in F, g \in G}$  indexed by  $F$  and  $G$ . Then from (11) it follows that

$$\dim Z_a \geq \text{rank } \langle (I - P_{\check{V}_a}) \bar{\Phi}_a, \bar{\Phi}_{a_+} \rangle.$$

We remark that if one finds a  $Z_a$  that satisfies (10), then one can always choose  $Z'_a \subset Z_a$  such that equality holds in the above estimate.

In this paper, we focus on constructions in which the spaces  $Z_a$  above are chosen via a generalization of the **intertwining technique** developed in [9] (and extended in [1] and [11]) to construct orthogonal piecewise polynomial wavelets. To that end, suppose  $\Phi'$  and  $\Phi''$  are bases centered on  $\mathbf{a}$  such that

$$(12) \quad S(\Phi) \subset S(\Phi') \subset S(\Phi'')$$

and that there exist spaces  $Z_a \subset \text{span } \check{\Phi}'_a$  (respectively,  $Z'_a \subset \text{span } \check{\Phi}''_a$ ),  $a \in \mathbf{a}$ , as in the previous paragraph, so that there exists an orthogonal basis  $\Omega$  (respectively,  $\Omega'$ ) such that  $S(\Omega) = S(\Phi) + S(Z)$  (respectively,  $S(\Omega') = S(\Phi') + S(Z')$ ). It then follows that

$$S(\Omega) \subset S(\Phi') \subset S(\Omega').$$

Specifically, suppose  $\mathbf{b}$  and  $\mathbf{c}$  are knot sequences in  $J$  so that  $\mathbf{a} \subset \mathbf{b} \subset \mathbf{c}$  and consider the spline bases  $\Phi = \Phi_{r,d}^{\mathbf{a}}$ ,  $\Phi' = \Phi_{r,d}^{\mathbf{b}}$ , and  $\Phi'' = \Phi_{r,d}^{\mathbf{c}}$  where  $r \geq 2d + 1$ . Then  $S(\Phi) \subset S(\Phi') \subset S(\Phi'')$ . We also recall that  $\Phi'$  and  $\Phi''$  are bases centered on  $\mathbf{a}$  as well as centered on  $\mathbf{b}$  and  $\mathbf{c}$ , respectively; see the remark following Lemma 1.

**1.3. Continuous, orthogonal, spline basis centered on a knot sequence: arbitrary polynomial reproduction.** In this example we will use extensively the results of [12]. Let  $J$  be an interval in  $\mathbf{R}$ ,  $\mathbf{a}$  a knot sequence in  $J$ , and let  $S_n^0(\mathbf{a})$  denote the spline space consisting of continuous piecewise polynomial functions in  $L^2(J)$  of degree at most  $n$  defined on  $J$  with break points in  $\mathbf{a}$ . Then  $S_n^0(\mathbf{a}) = S(\Phi)$  for a basis  $\Phi$  centered on  $\mathbf{a}$ . Let  $I = [0, 1]$  and  $\Pi^n$  denote the space of polynomials of degree  $n$  on  $I$ . If we set  $\tilde{\phi}^i(x) = x(1-x)p_{i-2}^{\frac{5}{2}}(2x-1)\chi_{[0,1)}(x)$ ,  $i = 2, 3, \dots, n$

where  $p_i^{\frac{5}{2}}(x)$  is the monic ultraspherical polynomial of degree  $i$  in  $x$  then  $\{\tilde{\phi}^i\}_{i=2}^n$  form an orthogonal set in  $L^2(I)$ . Let  $r = x\chi_{[0,1)}(x)$ ,  $l = (1-x)\chi_{[0,1)}(x)$ ,

$$(13) \quad U_n = \text{span} \{\tilde{\phi}^k \mid k = 2, \dots, n\},$$

$$(14) \quad r_n = (I - P_{U_n})r, \text{ and } l_n = (I - P_{U_n})l.$$

Some useful integrals that we will need later are

$$(15) \quad \|\tilde{\phi}^n\|^2 = \frac{(n-2)!(n+2)!}{16(2n-1)!!(2n+1)!!},$$

$$(16) \quad \langle r, \tilde{\phi}^n \rangle = \frac{(n-2)!}{4(2n-1)!!}, \quad \langle l, \tilde{\phi}^n \rangle = (-1)^n \langle r, \tilde{\phi}^n \rangle,$$

and

$$(17) \quad \langle r_n, l_n \rangle = (-1)^{n+1} \frac{1}{n(n+1)(n+2)}.$$

Here we use the notation that  $(2n-1)!! = (2n-1)(2n-3)\cdots(1)$ . These formulas may be obtained from [12] (cf., Equation 2.3 with  $m = 2$ ; Equation 2.5 with  $i = 0$ ,  $k = 0$ ; and Equation 5.3 where  $\tilde{\phi}^n(x) = \frac{1}{4}\phi_n^0(2x-1)\chi_{[0,1)}(x)$ ,  $r(x) = \frac{1}{2}r_0^0(2x-1)\chi_{[0,1)}(x)$  and  $l(x) = \frac{1}{2}l_0^0(2x-1)\chi_{[0,1)}(x)$ ). Since  $r_n$  and  $l_n$  are not orthogonal we add a function  $z^n$  chosen so that when it is projected out from the above two functions they become orthogonal. To accomplish this set,

$$(18) \quad z^n = \alpha_n \tilde{\phi}^{n+1} + \tilde{\phi}^{n+3},$$

where  $\alpha_n$  is fixed so that

$$(19) \quad \|z^n\|^2 \langle r_n, (I - P_{z^n})l_n \rangle = \langle r_n, l_n \rangle \|z^n\|^2 - (-1)^{n+1} \langle r, z^n \rangle^2 = 0,$$

where (16) has been used to obtain the above equation. The choice of  $z^n$  above is for simplicity and to preserve the symmetry properties of the basis being constructed. Substituting (18) into (19) and using the integrals computed above yields the quadratic equation

$$\alpha_n^2 + \left( \frac{2(n+1)}{2n+5} \right) \alpha_n + \left( \frac{(n+2)(n+1)(n^2-5n-30)}{(2n+7)((2n+5)^2(2n+3))} \right) = 0.$$

Choosing the positive square root in the quadratic formula we obtain the solution

$$(20) \quad \alpha_n = -\frac{n+1}{2n+5} + \frac{n+3}{2n+5} \sqrt{\frac{3(n+1)(n+3)}{(2n+7)(2n+3)}}.$$

Now we set

$$r^n = (I - P_{\{\tilde{\phi}^2, \dots, \tilde{\phi}^n, z^n\}})r = (I - P_{\{z^n\}})r_n$$

and

$$l^n = (I - P_{\{\tilde{\phi}^2, \dots, \tilde{\phi}^n, z^n\}})l = (I - P_{\{z^n\}})l_n.$$

Let  $\tilde{\Pi}^n = \text{span}\{\Pi^n, z^n\}$ . The above construction shows,

**Lemma 3.** *The functions  $\{r^n, l^n, z^n, \tilde{\phi}^i \mid i = 2, \dots, n\}$  form an orthogonal basis for  $\tilde{\Pi}^n$ . Furthermore  $\tilde{\Pi}^n \subset \tilde{\Pi}^{n+3}$ .*

We now construct a continuous orthogonal basis centered on  $\mathbf{a}$ . Let  $\sigma_a$  be the affine function taking  $a$  to 0 and  $a_+$  to 1. For each  $a < \sup J$  let  $\tilde{\phi}_a^n = \tilde{\phi}^n \circ \sigma_a$ ,  $z_a^n = z^n \circ \sigma_a$ ,  $r_a^n = r^n \circ \sigma_a$ , and  $l_a^n = l^n \circ \sigma_a$ . For  $a < \sup J$  define  $\check{\Omega}_a^n$  as

$$\check{\Omega}_a^n = \begin{cases} \{l_a^n, z_a^n, \tilde{\phi}_a^2, \dots, \tilde{\phi}_a^n\} & \text{if } a = \inf J \\ \{z_a^n, \tilde{\phi}_a^2, \dots, \tilde{\phi}_a^n\} & \text{if } \inf J < a < a_+ < \sup J \\ \{r_a^n, z_a^n, \tilde{\phi}_a^2, \dots, \tilde{\phi}_a^n\} & \text{if } a_+ = \sup J \end{cases}.$$

and, for  $\inf J < a < \sup J$ ,

$$\bar{\Omega}_a^n = \{r_{a-}^n + l_a^n\}.$$

From the above construction we see that  $\check{\Omega}_a^n$  and  $\bar{\Omega}_a^n$ , for  $a \in \mathbf{a}$ , form an orthogonal basis  $\Omega_{\mathbf{a}}^n$  centered on the knot sequence  $\mathbf{a}$  and that

$$(21) \quad S_n^0(\mathbf{a}) \subset S(\Omega_{\mathbf{a}}^n) \subset S_{n+3}^0(\mathbf{a}) \subset S(\Omega_{\mathbf{a}}^{n+3})$$

which implies

**Proposition 4.** *Let  $\mathbf{a}$  be a knot sequence and, for a fixed  $\kappa \in \{1, 2, 3\}$ , let  $V^k := S(\Omega_{\mathbf{a}}^{3k+\kappa})$ ,  $k = 0, 1, 2, \dots$ . Then  $V^k \subset V^{k+1}$  and  $\bigcup_{k=0}^{\infty} V^k$  is dense in  $L^2(\mathbf{J})$ .*

*Proof.* The first two statements follow from equation (21). The same equation also implies that  $\bigcup_{k=0}^{\infty} S_{3k+\kappa}^0(\mathbf{a}) \subset \bigcup_{k=0}^{\infty} V^k$ . The third statement now follows by the density of the piecewise continuous polynomials with breakpoints on  $\mathbf{a}$  in  $L^2(\mathbf{J})$ .  $\square$

**1.4. Continuous, orthogonal, spline bases centered on nested knot sequences.** Let  $J$  be an interval in  $\mathbf{R}$ ,  $\mathbf{a}$  a knot sequence in  $J$ , and let  $S_2^0(\mathbf{a})$  denote the spline space consisting of continuous piecewise quadratic spline functions defined on  $J$  with break points in  $\mathbf{a}$ . Then  $S_2^0(\mathbf{a}) = S(\Phi)$  for a basis  $\Phi$  centered on  $\mathbf{a}$ . For  $a < \sup J$ , we define the piecewise quadratic function  $\check{\phi}_a(x) = (x - a)(a_+ - x)\chi_{[a, a_+]}(x)$ , where  $\chi_A$  denotes the characteristic function of a set  $A \subset \mathbf{R}$ . Then for  $a < \sup J$ ,

$$\check{V}_a = \begin{cases} \text{span} \left\{ \check{\phi}_a, \left( \frac{a_+ - \cdot}{a_+ - a} \right) \chi_{[a, a_+]} \right\} & \text{if } a = \inf J \\ \text{span} \{ \check{\phi}_a \} & \text{if } \inf J < a < a_+ < \sup J \\ \text{span} \left\{ \check{\phi}_a, \left( \frac{\cdot - a}{a_+ - a} \right) \chi_{[a, a_+]} \right\} & \text{if } a_+ = \sup J \end{cases}.$$

Also, for  $\inf J < a < \sup J$ ,  $\bar{\Phi}_a$  can be chosen to consist of the piecewise linear spline  $\bar{\phi}_a$  that is 1 at  $a$  and 0 at  $b$  for  $b \neq a$ ,  $b \in \mathbf{a}$ .

We first construct a collection of orthogonal functions supported on the interval  $[0, 1)$ . These functions will then be used to construct an orthogonal continuous piecewise quadratic basis centered on the knot sequence  $\mathbf{a}$ . This construction was first given in [11] (we warn the reader that there are some typographical errors in the intermediate computations given in that paper that we now take the opportunity to correct).

Let  $r$  and  $l$  be as in the previous example and let  $q := 4rl$  denote the “quadratic bump” function of height one on  $[0, 1)$ . For  $0 < \theta < 1$ , let

$$q_0^\theta(x) := q(x/\theta), \quad q_1^\theta(x) := q((x - \theta)/(1 - \theta)),$$

and

$$h^\theta(x) = r(x/\theta) + l((x - \theta)/(1 - \theta)).$$

Now we look for a function  $z^\theta \in \text{span}\{q_0^\theta, q_1^\theta, h^\theta\}$  so that the

$$(22) \quad (I - P_{\text{span}\{q, z^\theta\}}) r \perp l$$

and  $z^\theta \perp q$ .

Note that  $q$  is in the 3-dimensional space  $\text{span}\{q_0^\theta, q_1^\theta, h^\theta\}$ . A basis for the 2-dimensional orthogonal complement of  $q$  in this space is given by (with help from *Mathematica*)

$$\begin{aligned} u_0 &= (1 - \theta)^2(2 + 3\theta)q_0^\theta + \theta^2(3\theta - 5)q_1^\theta \\ u_1 &= (-2 + 3(-1 + \theta)\theta^3)q_0^\theta + (-2 + 3(-1 + \theta)^3\theta)q_1^\theta + \left(\frac{16}{5} - 12(-1 + \theta)^2\theta^2\right)h^\theta. \end{aligned}$$

Then  $z^\theta$  must be of the form  $z^\theta = c_0u_0 + c_1u_1$ . Again using *Mathematica* we find that condition (22) is equivalent to the following quadratic equation in the variable  $c = c_0/c_1$

$$(23) \quad 0 = 5 \left( 4 - 5(1 - \theta)^2\theta^2(15 + (1 - \theta)\theta) \right) - 20(2 + \theta(9 + 13\theta(-3 + 2\theta)))c + 4(1 + 45(1 - \theta)\theta)c^2.$$

The discriminant of this equation is  $80 \left( 4 - 15(1 - \theta)^2\theta^2 \right)^2$  and thus is strictly positive for  $\theta \in (0, 1)$  giving the two solutions

$$c = \frac{20(2 + \theta(9 + 13\theta(2\theta - 3))) \pm 4\sqrt{5}(4 - 15(1 - \theta)^2\theta^2)}{8(1 + 45(1 - \theta)\theta)}.$$

Hence, for  $0 < \theta < 1$ , there is some  $z^\theta$  in the span of  $u_0$  and  $u_1$  such that (22) holds. Let  $r^\theta = (I - P_{\text{span}\{q, z^\theta\}})r$  and  $l^\theta = (I - P_{\text{span}\{q, z^\theta\}})l$ . Then  $\{r^\theta, l^\theta, q, z^\theta\}$  is an orthogonal system spanning a four dimensional space of piecewise quadratic functions continuous on  $[0, 1)$ .

We are now prepared to construct a continuous orthogonal basis centered on  $\mathbf{a}$  using the functions  $\{r^\theta, l^\theta, q, z^\theta\}$ . For  $a \in \mathbf{a}$  and  $a < \sup J$ , let  $\theta_a$  be in  $(0, 1)$  and let  $\boldsymbol{\theta}$  denote the sequence  $(\theta_a)_{a < \sup J}$ . For each  $a < \sup J$ , let  $z_a := z^{\theta_a} \circ \sigma_a$  where  $\sigma_a$  is the affine function taking  $a$  to 0 and  $a_+$  to 1. Let  $Z = Z_{\mathbf{a}, \boldsymbol{\theta}} := \Sigma_{a < \sup J} \text{span}\{z_a\}$ , then  $V = S_2^0(a) + Z$  satisfies the hypotheses of Lemma 2 and so  $V = S(\Omega)$  for some orthogonal basis  $\Omega = \Omega_{\mathbf{a}, \boldsymbol{\theta}}$  centered on  $\mathbf{a}$ . Specifically,  $\Omega$  can be chosen as follows: For  $a < \sup J$ ,

$$\check{\Omega}_a = \begin{cases} \{l^{\theta_a} \circ \sigma_a, q \circ \sigma_a, z_a\} & \text{if } a = \inf J \\ \{q \circ \sigma_a, z_a\} & \text{if } \inf J < a < a_+ < \sup J \\ \{r^{\theta_a} \circ \sigma_a, q \circ \sigma_a, z_a\} & \text{if } a_+ = \sup J \end{cases}.$$

and, for  $\inf J < a < \sup J$ ,  $\bar{\Omega}_a = \{r^{\theta_{a-}} \circ \sigma_{a-} + l^{\theta_a} \circ \sigma_a\}$ . Let  $\mathbf{b} = \mathbf{b}(\mathbf{a}, \boldsymbol{\theta})$  denote the sequence with elements  $b_a := (1 - \theta_a)a + \theta_a a_+$  between  $a$  and  $a_+$  for  $a < \sup J$ . Then

$$(24) \quad S_2^0(\mathbf{a}) \subset S(\Omega_{\mathbf{a}, \boldsymbol{\theta}}) \subset S_2^0(\mathbf{a} \cup \mathbf{b}).$$

**Lemma 5.** Let  $\mathbf{a}^0 \subset \mathbf{a}^1$  be knot sequences in  $J$ , let  $\boldsymbol{\theta}^0$  (indexed by  $\mathbf{a}^0$ ) and  $\boldsymbol{\theta}^1$  (indexed by  $\mathbf{a}^1$ ) be parameter sequences taking values in  $(0, 1)$ . For  $a \in \mathbf{a}^0$  let  $a_+$  denote the successor to  $a$  in  $\mathbf{a}^0$ . Let  $\mathbf{b}^0 := \mathbf{b}(\mathbf{a}^0, \boldsymbol{\theta}^0)$ ,  $\mathbf{b}^1 := \mathbf{b}(\mathbf{a}^1, \boldsymbol{\theta}^1)$  be such that  $b_a^0 \in \mathbf{a}^1$  whenever  $(a, a_+)$  contains some point in  $\mathbf{a}^1$ , and  $b_a^0 \in \mathbf{b}^1$  whenever  $(a, a_+)$  contains no point of  $\mathbf{a}^1$ .

Then

$$S(\Omega_{\mathbf{a}^0, \boldsymbol{\theta}^0}) \subset S(\Omega_{\mathbf{a}^1, \boldsymbol{\theta}^1}).$$

*Proof.* Let  $Z^\epsilon = Z_{\mathbf{a}^\epsilon, \theta^\epsilon}$  for  $\epsilon = 0, 1$ . Then  $S(\Omega_{\mathbf{a}^\epsilon, \theta^\epsilon}) = S_0^2(\mathbf{a}^\epsilon) + Z^\epsilon$ . Since  $S_0^2(\mathbf{a}^0) \subset S_0^2(\mathbf{a}^1)$  it suffices to show that  $Z^0 \subset S_0^2(\mathbf{a}^1) + Z^1$ . Let  $a \in \mathbf{a}^0$  with  $a < \sup J$ . If  $(a, a_+)$  contains no point of  $\mathbf{a}^1$  then  $a_+$  is also the successor of  $a$  in  $\mathbf{a}^1$ . Otherwise,  $b_a^0 \in \mathbf{a}^1$  in which case  $z_a^0 \in S(\mathbf{a}^1)$  and so, in either case, we have  $z_a^0 \in S_0^2(\mathbf{a}^1) + Z^1$  which completes the proof.  $\square$

**Proposition 6.** *Suppose  $(\mathbf{a}^k)_{k \in \mathbf{Z}}$  is a sequence of nested knot sequences in an interval  $J$  and  $(\theta^k)_{k \in \mathbf{Z}}$  is a sequence of parameter sequences taking values in  $(0, 1)$  are such that for each  $k \in \mathbf{Z}$ ,  $(\mathbf{a}^k, \theta^k)$  and  $(\mathbf{a}^{k+1}, \theta^{k+1})$  satisfy the hypotheses of Lemma 5. Let  $V^k := S(\Omega_{\mathbf{a}^k, \theta^k})$ . The following statements hold.*

- (1)  $V^k \subset V^{k+1}$  for  $(k \in \mathbf{Z})$ .
- (2) If  $\bigcup_k \mathbf{a}^k$  is dense in  $J$ , then  $\bigcup_k V^k$  is dense in  $L^2(J)$ .
- (3) Let  $\alpha := \bigcap_k \mathbf{a}^k \cup \mathbf{b}(a^k, \theta^k)$ . Then  $\bigcap_k V^k \subset S_2^0(\alpha)$ .

*Remark 2.* In the above,  $\alpha$  may not be a knot sequence in  $J$ , i.e., it may be empty, a singleton, or it may be that the  $\sup \alpha < \sup J$  or  $\inf \alpha > \inf J$ . In these cases we define  $S_2^0(\alpha)$  to be the collection of continuous functions on  $J$  whose restriction to any interval in  $J \setminus \alpha$  is in  $\Pi^d$ . For example  $S_2^0(\emptyset) = \Pi^d$ . We also mention that if  $\bigcup_k \mathbf{a}^k$  is dense in  $J$ , then  $\mathbf{b}(a^k, \theta^k) \subset \mathbf{a}^\ell$  for some  $\ell$  and so  $\alpha := \bigcap_k \mathbf{a}^k$  in this case.

*Proof.* Part (1) follows directly from Lemma 5. Equation (24) gives

$$S_2^0(\mathbf{a}^k) \subset V_k \subset S_2^0(\mathbf{a}^k \cup \mathbf{b}(a^k, \theta^k))$$

which shows (a)  $\bigcup_k S_2^0(\mathbf{a}^k) \subset \bigcup_k V^k$  and (b)  $\bigcap_k V^k \subset \bigcap_k S_2^0(\mathbf{a}^k \cup \mathbf{b}(a^k, \theta^k)) = S_2^0(\alpha)$  which implies parts (2) and (3).  $\square$

*Remark 3.* With  $V^k$  as in Lemma 6, let  $W^k$  denote the orthogonal complement of  $V^k$  in  $V^{k+1}$  and we have the usual decomposition

$$L^2(J) = \left( \bigcap_k V^k \right) \oplus \bigoplus_k W^k.$$

## 2. WAVELETS

In this section we consider nested spaces  $V^0 \subset V^1$  generated by orthogonal bases  $\Phi^0$  and  $\Phi^1$ , respectively, that are centered on the same knot sequence  $\mathbf{a}$ . For example, if  $(\mathbf{a}^0, \mathbf{b}^0)$  and  $(\mathbf{a}^1, \mathbf{b}^1)$  satisfy the conditions of Lemma 5, then  $V^0 = S(\Omega_{\mathbf{a}^0, \mathbf{b}^0})$  and  $V^1 = S(\Omega_{\mathbf{a}^1, \mathbf{b}^1})$  are such spaces centered on the *same* knot sequence  $\mathbf{a}^0$ . A second example, for  $n \geq 2$ , is given by  $V^0 = S(\Omega_{\mathbf{a}}^n)$  and  $V^1 = S(\Omega_{\mathbf{a}}^{n+3})$  where  $\Omega_{\mathbf{a}}^n$  is as in Section 1.3.

Our main result is that the **wavelet space**  $W := V^1 \ominus V^0$  has an orthogonal basis  $\Psi$  centered on  $\mathbf{a}$ .

**Theorem 7.** *Suppose  $\Phi^0$  and  $\Phi^1$  are orthogonal bases centered on a common knot sequence  $\mathbf{a}$  such that  $V^0 \subset V^1$  where  $V^0 = S(\Phi^0)$  and  $V^1 = S(\Phi^1)$ . Then there exists an orthogonal basis  $\Psi$  centered on  $\mathbf{a}$  such that  $W = S(\Psi)$  satisfies  $V^1 = V^0 \oplus W$ .*

*Proof.* The spaces  $S(\Phi^0)$  and  $S(\Phi^1)$  satisfy the conditions in Lemmas 1 and 2. Let  $V^0 := S(\Phi^0)$  and  $V^1 := S(\Phi^1)$ . For each  $a \in \mathbf{a}$ , define

$$\bar{V}_a^\epsilon := \left( I - P_{\check{V}_{a-}^\epsilon \oplus \check{V}_a^\epsilon} \right) V_a^\epsilon, \quad \epsilon = 0, 1$$

as in Lemma 2. It follows from (8) that  $\bar{V}_a^\epsilon \perp \bar{V}_{a+}^\epsilon$  and  $V_a^\epsilon = \check{V}_{a-}^\epsilon \oplus \bar{V}_a^\epsilon \oplus \check{V}_a^\epsilon$  for all  $a \in \mathbf{a}$  and  $\epsilon = 0, 1$ . Letting  $\bar{V}^\epsilon := S(\bar{\Phi}^\epsilon)$  and  $\check{V}^\epsilon := S(\check{\Phi}^\epsilon)$ , we have  $V^\epsilon = \bar{V}^\epsilon \oplus \check{V}^\epsilon$  for  $\epsilon = 0, 1$ .

Let  $W := V^1 \ominus V^0$ . First we describe the spaces  $\check{W}_a := W_{[a, a+]} = \check{V}_a^1 \cap (V^0)^\perp$  and  $W_a := W_{[a-, a+]}$  for  $a \in \mathbf{a}$ . Then we verify that  $W$  satisfies the hypotheses of Lemmas 1 and 2.

For  $a \in \mathbf{a}$ , observe that  $\bar{V}_a^0 \subset V_a^0 \subset V_a^1 = \check{V}_{a-}^1 \oplus \bar{V}_a^1 \oplus \check{V}_a^1$ , and thus

$$(25) \quad \bar{V}_a^0 \subset P_{\check{V}_{a-}^1} \bar{V}_a^0 \oplus P_{\bar{V}_a^1} \bar{V}_a^0 \oplus P_{\check{V}_a^1} \bar{V}_a^0 \subset A_a^- \oplus \bar{V}_a^1 \oplus A_a^+$$

where, for  $a \in \mathbf{a}$ ,

$$(26) \quad A_a^+ := P_{\check{V}_a^1} \bar{V}_a^0 \text{ and } A_a^- := P_{\check{V}_{a-}^1} \bar{V}_a^0.$$

**Lemma 8.** *For  $a \in \mathbf{a}$ , the spaces  $\check{V}_a^0$ ,  $A_a^+$ , and  $A_{a+}^-$  are mutually orthogonal subspaces of  $\check{V}_a^1$  and  $P_{\check{V}_a^1} V^0 = \check{V}_a^0 \oplus A_a^+ \oplus A_{a+}^-$ .*

*Proof.* Since  $\bar{V}_a^0, \bar{V}_{a+}^0 \perp \check{V}_a^0$  and  $\check{V}_a^0 \subset \check{V}_a^1$ , we have  $\bar{V}_a^0, \bar{V}_{a+}^0 \perp P_{\check{V}_a^1} \check{V}_a^0$ . Hence,  $A_a^+ := P_{\check{V}_a^1} \bar{V}_a^0$  and  $A_{a+}^- := P_{\check{V}_a^1} \bar{V}_{a+}^0$  are both perpendicular to  $\check{V}_a^0$  (note that if  $U_0$  and  $U_1$  are subspaces in a Hilbert space and  $P$  is an orthogonal projection such that  $U_0 \perp PU_1$  then  $PU_0 \perp U_1$ ). Let  $f \in \bar{V}_a^0$ . By Equation (25) we have  $f = P_{\check{V}_{a-}^1} f + P_{\bar{V}_a^1} f + P_{\check{V}_a^1} f$ . Similarly, if  $g \in \bar{V}_{a+}^0$ , then  $g = P_{\check{V}_a^1} g + P_{\bar{V}_{a+}^1} g + P_{\check{V}_{a+}^1} g$ . Hence, we have  $0 = \langle f, g \rangle = \langle P_{\check{V}_a^1} f, P_{\check{V}_a^1} g \rangle$  by the orthogonality assumption of  $V^1$ . Hence,  $A_a^+ \perp A_{a+}^-$ . The final equation follows from  $P_{\check{V}_a^1} V^0 = P_{\check{V}_a^1} (\check{V}_a^0 \oplus \bar{V}_{a-}^1 \oplus \bar{V}_a^1)$ .  $\square$

From Lemma 8 we obtain

$$(27) \quad \check{W}_a = \check{V}_a^1 \cap (V^0)^\perp = \check{V}_a^1 \cap (P_{\check{V}_a^1} V^0)^\perp = \check{V}_a^1 \cap (\check{V}_a^0 \oplus A_a^+ \oplus A_{a+}^-)^\perp,$$

and so

$$(28) \quad \check{V}_a^1 = \check{V}_a^0 \oplus A_a^+ \oplus A_{a+}^- \oplus \check{W}_a.$$

Then, again using Lemma 8, we have

$$(29) \quad \begin{aligned} (I - P_{V^0}) \check{V}_a^1 &\subset (I - P_{V^0}) A_a^+ + (I - P_{V^0}) A_{a+}^- + \check{W}_a \\ &= (I - P_{\bar{V}_a^0}) A_a^+ + (I - P_{\bar{V}_{a+}^0}) A_{a+}^- + \check{W}_a \\ &\subset W_a + W_{a+} \end{aligned}$$

where we observe that  $(I - P_{\bar{V}_a^0}) A_a^+ \subset (\bar{V}_a^0 + A_a^+) \cap V_0^\perp \subset W_a$  and similarly  $(I - P_{\bar{V}_{a+}^0}) A_{a+}^- \subset W_{a+}$ . Furthermore,

$$(30) \quad (I - P_{V^0}) \bar{V}_a^1 = (I - P_{\bar{V}_a^0}) \bar{V}_a^1 \subset W_a,$$

and, hence,

$$(31) \quad (I - P_{V^0}) V_a^1 \subset W_{a-} + W_a + W_{a+}.$$

Letting  $\Sigma_{a \in \mathbf{a}}^* V_a$  denote  $\text{clos}_{L^2(J)} \text{span} \left( \bigcup_{a \in \mathbf{a}} V_a \right)$  for a space  $V$  centered on  $\mathbf{a}$ , we have

$$(32) \quad W = (I - P_{V^0}) \Sigma_{a \in \mathbf{a}}^* V_a^1 = \Sigma_{a \in \mathbf{a}}^* (I - P_{V^0}) V_a^1 = \Sigma_{a \in \mathbf{a}}^* W_a,$$

that is,  $W$  satisfies  $(\hat{b})$  of Lemma 1.

If  $f \in W$  vanishes on  $[a, a_+]$ , then  $f = f_1 + f_2$  where  $f_1 \in V_{(-\infty, a]}^1$  and  $f_2 \in V_{[a_+, \infty)}^1$  (property  $(\hat{c})$  of Lemma 1 applied to  $V^1$ ). Since  $f \perp V^0$  and using the support properties of  $f_1$  and  $f_2$ , we have

$$0 = P_{V^0} f = P_{V^0} f_1 + P_{V^0} f_2 = P_{\bar{V}_a^0} f_1 + P_{\bar{V}_{a_+}^0} f_2.$$

Since  $\bar{V}_a^0 \perp \bar{V}_{a_+}^0$  we must have  $P_{\bar{V}_a^0} f_1 = P_{\bar{V}_{a_+}^0} f_2 = 0$  and hence  $f_1 \in W_{(-\infty, a]}$  and  $f_2 \in W_{[a_+, \infty)}$  which verifies that  $W$  satisfies property  $(\hat{c})$  of Lemma 1. Since  $W_a \subset V_a^1$  is finite dimensional, then Lemma 1 implies  $W = S(\Psi)$  for some basis  $\Psi$  centered on  $\mathbf{a}$ .

Observe that

$$\begin{aligned} (33) \quad W_a &\subset V_a^1 \cap (\check{V}_a^0 \oplus \bar{V}_{a_+}^0)^\perp \subset \check{V}_{a_-}^1 \oplus \bar{V}_a^1 \oplus (\check{V}_a^1 \cap (\check{V}_a^0 \oplus \bar{V}_{a_+}^0))^\perp \\ &= \check{V}_{a_-}^1 \oplus \bar{V}_a^1 \oplus A_a^+ \oplus \check{W}_a, \end{aligned}$$

and in the same way we have

$$(34) \quad W_{a_+} \subset A_{a_+}^- \oplus \check{W}_a \oplus \check{V}_{a_+}^1 \oplus \bar{V}_{a_+}^1.$$

It then follows that  $(I - P_{\check{W}_a})W_a \perp W_{a_+}$  and so by Lemma 2 we conclude that  $W = S(\Psi)$  for some orthogonal basis centered on  $\mathbf{a}$ .

We recall that in the case that  $J$  is not an open interval and  $a \in \mathbf{a}$  is an endpoint of  $J$ , say  $a = \inf J$ , that  $a_- = -\infty$  and that the spaces  $\check{V}_{a_-}^\epsilon$ ,  $\bar{V}_a^\epsilon$  are trivial for  $\epsilon = 0, 1$ . It then follows that  $\bar{W}_a = \{0\}$  whenever  $a$  is an endpoint of  $J$ .  $\square$

The spaces  $A_a^\pm$  defined in (26) play central roles in the above proof. In the following corollary, we express  $\check{W}_a$  and  $\bar{W}_a$  in terms of these spaces.

**Corollary 9.** *Suppose  $V^0$ ,  $V^1$ , and  $W$  are as in Theorem 7 and let  $A_a^\pm$  be given by (26). Let  $\bar{W}_a := W_a \ominus (\check{W}_{a_-} \oplus \check{W}_a)$ . Then*

$$(35) \quad \check{W}_a = \check{V}_a^1 \cap (\check{V}_a^0 \oplus A_a^+ \oplus A_{a_+}^-)^\perp,$$

and

$$(36) \quad \bar{W}_a = (A_a^- \oplus \bar{V}_a^1 \oplus A_a^+) \cap (\bar{V}_a^0)^\perp.$$

*Proof.* The formula (35) is given in (27). Proceeding as in the previous proof,

$$\begin{aligned} (37) \quad W_a &= V_a^1 \cap (V_{a_-}^0 \oplus \bar{V}_a^0 \oplus V_{a_+}^0)^\perp \\ &= (A_a^- \oplus \bar{V}_a^1 \oplus A_a^+ \oplus \check{W}_{a_-} \oplus \check{W}_a) \cap (\bar{V}_a^0)^\perp \\ &= ((A_a^- \oplus \bar{V}_a^1 \oplus A_a^+) \cap (\bar{V}_a^0)^\perp) \oplus \check{W}_{a_-} \oplus \check{W}_a \\ &= \bar{W}_a \oplus \check{W}_{a_-} \oplus \check{W}_a. \end{aligned}$$

which establishes (36).  $\square$

**2.1. Decomposing the wavelet spaces.** In order to aid in the construction of the wavelets, we discuss a parallel development based on the wavelet construction given in [12]. This construction decomposes  $\bar{W}_a$  into two orthogonal subspaces that illuminate and simplify the construction of the bar wavelets. We also give more explicit characterizations of the dimensions of the subspaces discussed above.

Let  $k_a^\epsilon = \dim V_a^\epsilon$ ,  $\bar{k}_a^\epsilon = \dim \bar{V}_a^\epsilon$ ,  $\check{k}_a^\epsilon = \dim \check{V}_a^\epsilon$ ,  $Y_a = \bar{V}_a^0 \cap \bar{V}_a^1$ ,  $m_a = \dim Y_a$ ,  $Y_{a-}^- = \bar{V}_a^0 \chi_{[a-,a]} \cap \bar{V}_a^1 \chi_{[a-,a]}$ ,  $\dim Y_{a-}^- = m_{a-}^-$ ,  $Y_{a+}^+ = \bar{V}_a^0 \chi_{[a,a+]} \cap \bar{V}_a^1 \chi_{[a,a+]}$ ,  $\dim Y_{a+}^+ = m_{a+}^+$ ,  $X_{a+}^+ = (\chi_{[a,a+]} \bar{V}_a^0) \ominus Y_{a+}^+$ , and  $X_{a+}^- := (\chi_{[a-,a]} \bar{V}_a^0) \ominus Y_{a-}^-$ .

We begin with a sort of algorithm for constructing the “short” wavelet space,  $\check{W}_a$ , which will show the use of the spaces,  $Y_{a+}^+$  and  $Y_{a-}^-$ . Recall that  $A_{a-}^- = P_{\check{V}_{a-}^1} \bar{V}_a^0$  and  $A_{a+}^+ = P_{\check{V}_a^1} \bar{V}_a^0$ .

**Lemma 10.** (1)  $\dim \check{W}_a = (\check{k}_a^1 - \check{k}_a^0) - \dim X_{a+}^+ - \dim X_{a+}^-$ .  
 (2)  $A_{a+}^+ = P_{\check{V}_a^1} X_{a+}^+$ , and  $A_{a+}^- = P_{\check{V}_a^1} X_{a+}^-$ .  
 (3)  $\dim A_{a+}^+ = \dim X_{a+}^+ = \bar{k}_a^0 - m_{a+}^+$ , and  $\dim A_{a+}^- = \dim X_{a+}^- = \bar{k}_{a+}^0 - m_{a+}^-$ .

*Proof.* From the definition of  $W$  it follows that  $\check{W}_a \subset \check{V}_a^1 \ominus \check{V}_a^0$ . But  $\check{W}_a$  is also orthogonal to  $\chi_{[a,a+]} \bar{V}_a^0 = X_{a+}^+ \oplus Y_{a+}^+$ , and  $\chi_{[a,a+]} \bar{V}_{a+}^0 = X_{a+}^- \oplus Y_{a+}^-$ . Since  $Y_{a+}^+$  and  $Y_{a+}^-$  are already orthogonal to  $\check{V}_a^1 \ominus \check{V}_a^0$ , then to construct a basis for  $\check{W}_a$  we only need to choose a basis from  $\check{V}_a^1 \ominus \check{V}_a^0$  that is simultaneously orthogonal to  $X_{a+}^+$  and  $X_{a+}^-$ . Further, since  $\check{V}_a^1 \ominus \check{V}_a^0 \subset \check{V}_a^1$ , we need this basis orthogonal to spaces  $P_{\check{V}_a^1} X_{a+}^+$  and  $P_{\check{V}_a^1} X_{a+}^-$ .

Using the fact that  $\bar{f}_a^0 = P_{\check{V}_{a-}^1} \bar{f}_a^0 + P_{\bar{V}_a^1} \bar{f}_a^0 + P_{\check{V}_a^1} \bar{f}_a^0$ , for  $\bar{f}_a^0 \in \bar{V}_a^0$ , it is easy to check that the kernel of  $P_{\check{V}_a^1}$  acting on  $\chi_{[a,a+]} \bar{V}_a^0$  is  $Y_{a+}^+$ . Thus  $\dim P_{\check{V}_a^1} X_{a+}^+ = \dim X_{a+}^+$ . Similarly it can be shown that  $\dim P_{\check{V}_a^1} X_{a+}^- = \dim X_{a+}^-$ . It is easy to check that the spaces  $P_{\check{V}_a^1} X_{a+}^+$  and  $P_{\check{V}_a^1} X_{a+}^-$  are subspaces of  $\check{V}_a^1 \ominus \check{V}_a^0$  and are orthogonal to each other. The first equation of the statement of the lemma follows now since the number of additional orthogonality conditions is  $\dim X_{a+}^+ + \dim X_{a+}^-$ .

Observe that  $A_{a+}^+ = P_{\check{V}_a^1} \chi_{[a,a+]} \bar{V}_a^0 = P_{\check{V}_a^1} (X_{a+}^+ \oplus Y_{a+}^+) = P_{\check{V}_a^1} X_{a+}^+$ . Similarly we can show that  $A_{a+}^- = P_{\check{V}_a^1} X_{a+}^-$ . By local linear independence,  $\dim \bar{V}_a^0 \chi_{[a,a+]} = \dim \bar{V}_a^0 = \bar{k}_a^0$  and so  $\dim A_{a+}^+ = \bar{k}_a^0 - m_{a+}^+$ . The formula for  $\dim A_{a+}^-$  follows in the same way.  $\square$

Using Lemma 10 combined with Corollary 9, we now have

**Theorem 11.**

$$\begin{aligned} \dim \bar{W}_a &= \bar{k}_a^1 + \bar{k}_a^0 - m_{a+}^+ - m_{a-}^-, \\ \dim \check{W}_a &= \check{k}_a^1 - \check{k}_a^0 - \bar{k}_a^0 - \bar{k}_{a+}^0 + m_{a+}^+ + m_{a+}^-, \\ \dim(\check{W}_{a-} \oplus \check{W}_a) &= (m_{a-}^+ - \bar{k}_{a-}^0) + (k_a^1 - k_a^0 - \bar{k}_a^1 - \bar{k}_a^0 + m_{a-}^- + m_{a+}^+) + (m_{a+}^- - \bar{k}_{a+}^0). \end{aligned}$$

We now introduce spaces that aid in the decomposition  $\bar{W}_a$ , this follows the development given in [12]. Let

$$\begin{aligned} T_a &= (I - P_{\bar{V}_a^1}) \bar{V}_a^0 \\ T_a^- &= \{f \in T_a : \text{supp } f \subset [a-, a]\} \\ T_a^+ &= \{f \in T_a : \text{supp } f \subset [a, a+]\} \\ U_a &= T_a \ominus (T_a^- \oplus T_a^+) \\ S_a &= (\chi_{[a,a+]} - \chi_{[a-,a]}) U_a \end{aligned}$$

*Remark 4.* In a typical refinement the spaces  $T_a^\pm$  are trivial. However in section 3.3 we present an example of a refinement where these spaces are not empty.

**Lemma 12.**  $A_a^- \oplus A_a^+ = S_a + T_a$ .

*Proof.* First we remark that

$$T_a = (P_{V_a^1} - P_{\check{V}_a^1})\bar{V}_a^0 = (P_{\check{V}_a^1} + P_{\check{V}_{a-}^1})\bar{V}_a^0 \subset P_{\check{V}_{a-}^1}\bar{V}_a^0 + P_{\check{V}_a^1}\bar{V}_a^0 = A_a^- \oplus A_a^+.$$

Next, define

$$\tilde{S}_a := (P_{\check{V}_a^1} - P_{\check{V}_{a-}^1})\bar{V}_a^0.$$

It is clear that

$$A_a^- \oplus A_a^+ = \tilde{S}_a + T_a.$$

Furthermore,

$$(P_{\check{V}_a^1} - P_{\check{V}_{a-}^1})T_a = (P_{\check{V}_a^1} - P_{\check{V}_{a-}^1})(P_{\check{V}_a^1} + P_{\check{V}_{a-}^1})\bar{V}_a^0 = (P_{\check{V}_a^1} - P_{\check{V}_{a-}^1})\bar{V}_a^0 = \tilde{S}_a$$

which shows

$$\tilde{S}_a = (P_{\check{V}_a^1} - P_{\check{V}_{a-}^1})T_a = (P_{\check{V}_a^1} - P_{\check{V}_{a-}^1})(U_a \oplus T_a^- \oplus T_a^+) = S_a + T_a^- + T_a^+;$$

where we used  $S_a = (\chi_{[a, a_+]} - \chi_{[a_-, a]})U_a = (P_{\check{V}_a^1} - P_{\check{V}_{a-}^1})U_a$  since  $U_a \subset A_a^- \oplus A_a^+$ .

Since  $T_a^-$  and  $T_a^+$  are contained in  $T_a$ , we have  $\tilde{S}_a + T_a = S_a + T_a$  which completes the proof.  $\square$

**Lemma 13.**  $\dim T_a = \bar{k}_a^0 - m_a$ ,  $\dim T_a^- = m_a^+ - m_a$ ,  $\dim T_a^+ = m_a^- - m_a$ ,  $\dim U_a = \bar{k}_a^0 + m_a - m_a^- - m_a^+ = \dim S_a$ .

*Remark 5.* We take this opportunity to point out that in [12, page 1041] the dimension of  $T_0$  should be  $m_1 - m$  and the dimension of  $T_1$  should be  $m_0 - m$ . The analog of these spaces are  $T_a^-$  and  $T_a^+$ .

*Proof.* The linear transformation  $I - P_{\check{V}_a^1}$  maps  $\bar{V}_a^0$  onto  $T_a$ . The kernel of this mapping is clearly  $Y_a$ , and thus the stated dimension of  $T_a$  follows.

Once the dimensions of  $T_a^+$  and  $T_a^-$ , as stated, are proved, the stated dimension of  $U_a$  follows easily from its definition. Since  $P_{\check{V}_a^1}U_a = 0$ ,

$$(38) \quad U_a \subset \check{V}_{a-}^1 \oplus \check{V}_a^1,$$

which implies that multiplying functions in  $U_a$  by characterisitic functions above does not destroy membership in  $V_a^1$  so the dimension of  $S_a$  follows.

Let  $y_a^+ \in Y_a^+$ . Then there are functions  $\bar{f}_a^0 \in \bar{V}_a^0$  and  $\bar{f}_a^1 \in \bar{V}_a^1$  so that  $y_a^+ = \bar{f}_a^0 \chi_{[a, a_+]} = \bar{f}_a^1 \chi_{[a, a_+]}$ . Note that by local linear independence the functions  $\bar{f}_a^0$  and  $\bar{f}_a^1$  are uniquely determined.

It follows that

$$(I - P_{\check{V}_a^1})\bar{f}_a^0 = P_{\check{V}_a^1}\bar{f}_a^0 + P_{\check{V}_a^1}\bar{f}_a^0 = P_{\check{V}_a^1}\bar{f}_a^0 + P_{\check{V}_a^1}\bar{f}_a^1 = P_{\check{V}_a^1}\bar{f}_a^0.$$

As such,  $(I - P_{\check{V}_a^1})\bar{f}_a^0 \in T_a^-$ .

Conversely, let  $t_a^- \in T_a^-$ . Choose  $\bar{f}_a^0 \in \bar{V}_a^0$  so that  $(I - P_{\check{V}_a^1})\bar{f}_a^0 = t_a^-$ . Since  $(I - P_{\check{V}_a^1})\bar{f}_a^0 = P_{\check{V}_a^1}\bar{f}_a^0 + P_{\check{V}_a^1}\bar{f}_a^0$  we get that  $P_{\check{V}_a^1}\bar{f}_a^0 = 0$ . By the same argument as given in the proof of Lemma 10 it follows that  $\bar{f}_a^0 \chi_{[a, a_+]} \in Y_a^+$ .

Define the linear transformation  $F : Y_a^+ \rightarrow T_a^-$  by  $F(y_a^+) = (I - P_{\bar{V}_a^1})\bar{f}_a^0$ . The above argument shows that  $F$  is onto. It is clear that the kernel of  $F$  is  $Y_a\chi_{[a, a_+]}$ . Since, by local linear independence,  $\dim \chi_{[a_-, a]}Y_a = \dim Y_a = m_a$ , it follows that  $\dim T_a^- = m_a^+ - m_a$ .

The proof that  $\dim T_a^+ = m_a^- - m_a$  is entirely analogous.  $\square$

*Remark 6.* From Lemmas 10, 12, and 13, it follows that  $\dim S_a + \dim T_a = \dim(S_a + T_a)$  and thus  $S_a$  and  $T_a$  are linearly independent.

We now begin the decomposition of  $\bar{W}_a$ .

**Lemma 14.** *Let  $\hat{W}_a := (I - P_{\bar{V}_a^0})\bar{V}_a^1$ . Then  $\dim \hat{W}_a = \bar{k}_a^1 - m_a$  and  $\hat{W}_a \subset \bar{W}_a$ .*

*Proof.* Since  $\bar{V}_a^0 \subset A_a^- \oplus \bar{V}_a^1 \oplus A_a^+$  it follows from Remark 2.2 that  $\hat{W}_a \subset \bar{W}_a$ . The dimension formula for  $\hat{W}_a$  follows from elementary Hilbert space arguments.  $\square$

**Lemma 15.**  $\bar{V}_a^0 \oplus \hat{W}_a = T_a \oplus \bar{V}_a^1$ .

*Proof.* Since the dimension of the spaces on each side of the above equations are equal we need only show that one space is contained in the other to prove the result. By definition of  $\hat{W}_a$  it follows that  $\bar{V}_a^1 \subset \bar{V}_a^0 \oplus \hat{W}_a$ . This, in turn, implies that  $T_a \subset \bar{V}_a^0 \oplus \hat{W}_a$ .  $\square$

**Lemma 16.** *Let  $\tilde{W}_a := \bar{W}_a \ominus \hat{W}_a$ . Then*

(1)

$$\begin{aligned} \tilde{W}_a &= (A_a^- \oplus A_a^+) \cap (\bar{V}_a^0)^\perp \\ &= (I - P_{\bar{V}_a^0 \oplus \hat{W}_a})(A_a^- \oplus A_a^+) \\ &= (I - P_{\bar{V}_a^0 \oplus \hat{W}_a})S_a. \end{aligned}$$

(2)  $\dim \tilde{W}_a = \bar{k}_a^0 + m_a - m_a^- - m_a^+$

*Proof.* (1) Let  $\tilde{w}_a \in \tilde{W}_a$ . Obviously,  $\tilde{w}_a \in (\bar{V}_a^0)^\perp$ . Choose  $f_{a_-, a} \in A_a^-$ ,  $\bar{f}_a^1 \in \bar{V}_a^1$ , and  $f_{a, a} \in A_a^+$  such that  $\tilde{w}_a = f_{a_-, a} + \bar{f}_a^1 + f_{a, a}$ . Now  $0 = \langle \tilde{w}_a, \bar{f}_a^1 - P_{\bar{V}_a^0}\bar{f}_a^1 \rangle = \langle \tilde{w}_a, \bar{f}_a^1 \rangle = \langle \bar{f}_a^1, \bar{f}_a^1 \rangle$ . Thus  $\bar{f}_a^1 = 0$  and  $\tilde{w}_a = f_{a_-, a} + f_{a, a}$ . We now have that  $\tilde{W}_a \subset (A_a^- \oplus A_a^+) \cap (\bar{V}_a^0)^\perp$ .

Next, choose  $f_{a_-, a} \in A_a^-$  and  $f_{a, a} \in A_a^+$  such that  $f_{a_-, a} + f_{a, a} \perp \bar{V}_a^0$ . By the definition of  $\hat{W}_a$ , it is clear that  $f_{a_-, a} + f_{a, a} \perp \hat{W}_a$ . Thus  $(I - P_{\bar{V}_a^0 \oplus \hat{W}_a})(f_{a_-, a} + f_{a, a}) = f_{a_-, a} + f_{a, a}$ . We then get that  $(A_a^- \oplus A_a^+) \cap (\bar{V}_a^0)^\perp \subset (I - P_{\bar{V}_a^0 \oplus \hat{W}_a})(A_a^- \oplus A_a^+)$ .

By the characterization of  $\bar{W}_a$  in the wavelet theorem, it is clear that  $(I - P_{\bar{V}_a^0 \oplus \hat{W}_a})(A_a^- \oplus A_a^+) \subset \bar{W}_a$ .

Now we have, by Lemmas 12 and 15,  $\tilde{W}_a = (I - P_{\bar{V}_a^0 \oplus \hat{W}_a})(A_a^- \oplus A_a^+) = (I - P_{\bar{V}_a^1 \oplus T_a})(S_a + T_a) = (I - P_{\bar{V}_a^1 \oplus T_a})S_a$ .

(2) This follows directly from Theorem 11 and Lemma 14.  $\square$

*Remark 7.* It follows from Lemmas 13 and 16 that  $(\bar{V}_a^0 \oplus \hat{W}_a) \cap S_a = 0$

**2.2. Algorithm.** The decomposition given in subsection 2.1 provides a procedure for constructing an orthonormal wavelet basis  $\Psi$  which we next summarize. Note that the construction is local and if the scaling functions are symmetric or antisymmetric on the interval then the wavelets can be constructed which are symmetric or antisymmetric.

Fix  $a \in \mathbf{a}$  and let  $k_a^\epsilon$ ,  $\bar{k}_a^\epsilon$ ,  $\check{k}_a^\epsilon$ ,  $m_a$ , and  $m_a^\pm$  be as in the first paragraph of subsection 2.1. We begin with the construction of the “long”  $\bar{W}_a$  wavelets. From Lemma 14

$$(39) \quad \hat{W}_a = (I - P_{\bar{V}_a^0})\bar{V}_a^1$$

which has dimension  $\bar{k}_a^1 - m_a$  so construct an orthonormal basis of  $\bar{k}_a^1 - m_a$  wavelet functions for this space. If the scaling functions have a symmetric or antisymmetric with respect to the point  $a$  then taking linear combinations of the symmetric functions or antisymmetric functions in  $\hat{W}_a$  allows us to construct wavelets functions that preserve the symmetry. Next construct  $T_a$ , then eliminate the functions in  $T_a$  which are supported in  $[a_-, a]$  or  $[a, a_+]$  to find  $U_a$ . Finally apply the difference in characteristic functions to obtain  $S_a$  as given below Theorem 11. Now by Gram-Schmidt construct an orthonormal basis of  $k_a^0 + m_a - m_a^- - m_a^+$  functions for

$$(40) \quad \tilde{W}_a = (I - P_{\bar{V}_a^0 \oplus \hat{W}_a})S_a$$

given by equation (1) of Lemma 16. From the construction of  $S_a$  and the discussion above we see that if the scaling functions are symmetric or antisymmetric with respect to  $a$  then the wavelets can be constructed to preserve the symmetry. The final step is to construct the wavelets in  $\tilde{W}_a$ . From equation (35) there are  $\check{k}_a^1 - \check{k}_a^0 - \bar{k}_a^0 - \bar{k}_{a_+}^0 + m_a^+ + m_{a_+}^-$  orthonormal functions to be computed in

$$(41) \quad \check{W}_a = (I - P_{\check{V}_a^0 \oplus A_a^+ \oplus A_{a_+}^-})\check{V}_a^1$$

to complete the construction of the wavelets.

**2.3. Scaling and Wavelet Matrix Coefficients .** In this section we need to make use of matrices consisting of inner products. Let  $f$  represent a column vector consisting of a finite collection of functions in  $L^2(\mathbf{R})$ . If  $g$  is another such column we let  $\langle f, g \rangle$  denote the matrix so that the  $ij$  term is given by  $\langle f_i, g_j \rangle$ .

Let  $\Phi^0$  and  $\Phi^1$  be orthonormal bases centered on a knot sequence  $\mathbf{a}$  as in Theorem 7, with associated orthonormal wavelet basis  $\Psi$ . For  $a' \in \{a_-, a\}$  define

$$\bar{c}_{aa'}^- := \langle \bar{\phi}_a^0, \bar{\phi}_{a'}^1 \rangle.$$

For  $a' \in \{a_-, a\}$  define

$$\tilde{c}_{aa'}^- := \langle \bar{\phi}_a^0, \check{\phi}_{a'}^1 \rangle.$$

Similarly, for  $a' \in \{a_-, a\}$ , we define  $\check{c}_{aa'}^-$  and  $\check{\check{c}}_{aa'}^-$ . (Note: Of course these matrices are only defined when the appropriate collections of functions are non-empty.)

It follows that

$$(42) \quad \begin{bmatrix} \check{\check{\Phi}}_{a_-}^0 \\ \check{\check{\Phi}}_a^0 \\ \check{\check{\Phi}}_a^0 \end{bmatrix} = \begin{bmatrix} \check{\check{c}}_{a_- a_-}^- & 0 & 0 \\ \check{\check{c}}_{aa_-}^- & \bar{c}_{aa}^- & \check{\check{c}}_{aa}^- \\ 0 & 0 & \check{\check{c}}_{aa}^- \end{bmatrix} \begin{bmatrix} \check{\check{\Phi}}_{a_-}^1 \\ \check{\check{\Phi}}_a^1 \\ \check{\check{\Phi}}_a^1 \end{bmatrix}$$

Note that in equation (42) we can regard the matrix

$$\begin{bmatrix} \check{c}_{a_- a_-} & 0 & 0 \\ \check{c}_{aa_-} & \check{c}_{aa} & \check{c}_{aa} \\ 0 & 0 & \check{c}_{aa} \end{bmatrix}$$

as having rows (resp. columns) indexed by

$$\begin{bmatrix} \check{\Phi}_{a_-}^0 \\ \check{\Phi}_a^0 \\ \check{\Phi}_a^0 \end{bmatrix} \left( \text{resp. } \begin{bmatrix} \check{\Phi}_{a_-}^1 \\ \check{\Phi}_a^1 \\ \check{\Phi}_a^1 \end{bmatrix} \right).$$

If one of the row (resp. column) indices is empty then the corresponding row (resp. column) is omitted. For example, if  $\check{\Phi}_a^0$  and  $\check{\Phi}_a^1$  are empty then equation (42) becomes

$$\begin{bmatrix} \check{\Phi}_{a_-}^0 \\ \check{\Phi}_a^0 \end{bmatrix} = \begin{bmatrix} \check{c}_{a_- a_-} & 0 \\ 0 & \check{c}_{aa} \end{bmatrix} \begin{bmatrix} \check{\Phi}_{a_-}^1 \\ \check{\Phi}_a^1 \end{bmatrix}.$$

We follow this convention in all the remaining matrix constructions.

For  $a' \in \{a_-, a\}$  we can also define matrices  $\check{d}_{aa'}$ ,  $\check{d}_{aa'}$ ,  $\check{d}_{aa'}$ , and  $\check{d}_{aa'}$  so that

$$(43) \quad \begin{bmatrix} \check{\Psi}_{a_-} \\ \check{\Psi}_a \\ \check{\Psi}_a \end{bmatrix} = \begin{bmatrix} \check{d}_{a_- a_-} & 0 & 0 \\ \check{d}_{aa_-} & \check{d}_{aa} & \check{d}_{aa} \\ 0 & 0 & \check{d}_{aa} \end{bmatrix} \begin{bmatrix} \check{\Phi}_{a_-}^1 \\ \check{\Phi}_a^1 \\ \check{\Phi}_a^1 \end{bmatrix}$$

For knot  $a \in \mathbf{a}$  and  $a' \in \{a_-, a\}$  define

$$c_{aa'} := \begin{bmatrix} \check{c}_{aa'} & \check{c}_{aa'} \\ \check{c}_{aa'} & \check{c}_{aa'} \end{bmatrix}.$$

In particular, we have

$$c_{aa_-} = \begin{bmatrix} 0 & \check{c}_{aa_-} \\ 0 & 0 \end{bmatrix} \text{ and } c_{aa} = \begin{bmatrix} \check{c}_{aa} & \check{c}_{aa} \\ 0 & \check{c}_{aa} \end{bmatrix}.$$

For  $a \in \mathbf{a}$ , equation (42) implies the following "scaling equation":

$$(44) \quad \Phi_a^0 = \sum_{a' \in \{a_-, a\}} c_{aa'} \Phi_{a'}^1.$$

Similarly we can define

$$d_{aa'} := \begin{bmatrix} \check{d}_{aa'} & \check{d}_{aa'} \\ \check{d}_{aa'} & \check{d}_{aa'} \end{bmatrix}.$$

For  $a \in \mathbf{a}$ , equation (43) implies

$$(45) \quad \Psi_a = \sum_{a' \in \{a_-, a\}} d_{aa'} \Phi_{a'}^1.$$

Below we describe the construction of  $\Psi$  in terms of the matrix coefficients  $d_{aa'}$ .

When they are defined, the matrices  $\check{c}_{aa}$  are full rank with orthonormal rows as is the block matrix

$$\begin{bmatrix} \check{c}_{aa_-} & \check{c}_{aa} & \check{c}_{aa} \end{bmatrix}.$$

However, the individual blocks may not be full rank (although this is the generic case). For  $a' = a, a_-$ , if  $\bar{c}_{aa'} \neq 0$  let  $\bar{b}_{aa'}$  be the matrix with orthonormal rows whose row span is the same as  $\bar{c}_{aa'}$  and let  $\bar{e}_{aa'}$  be the matrix such that

$$\bar{c}_{aa'} = \bar{e}_{aa'} \bar{b}_{aa'}.$$

We observe that

$$\alpha_a^+ := \bar{b}_{aa}^{\check{}} \check{\Phi}_a^1$$

is an orthonormal basis for  $A_a^+$ , and

$$\alpha_a^- := \bar{b}_{aa-}^{\check{}} \check{\Phi}_{a-}^1$$

is an orthonormal basis for  $A_a^-$ . (Note: If  $\bar{b}_{aa}^{\check{}}$  (resp.  $\bar{b}_{aa-}^{\check{}}$ ) is 0 then  $\alpha_a^+$  (resp.  $\alpha_a^-$ ) is empty.

From Corollary 9 we have

$$\check{W}_a := \check{V}_a^1 \ominus \left( \check{V}_a^0 \oplus A_a^+ \oplus A_{a+}^- \right).$$

If  $\check{W}_a \neq \{0\}$  it follows that  $\check{d}_{aa}^{\check{}}$  may be chosen so that

$$\begin{bmatrix} \bar{c}_{aa}^{\check{}} \\ \bar{b}_{aa}^{\check{}} \\ \bar{b}_{a+}^{\check{}} \\ \check{d}_{aa}^{\check{}} \end{bmatrix}$$

is a square orthogonal matrix which means that  $\check{\Psi}_a = \check{d}_{aa}^{\check{}} \check{\Phi}_a^1$ . If  $\check{W}_a = 0$ , then  $\check{\Psi}_a$  is empty.

From equation (42) we see that, when defined,

$$\begin{aligned} \bar{\Phi}_a^0 &= \begin{bmatrix} \bar{c}_{aa-}^{\check{}} & \bar{c}_{aa}^{\check{}} & \bar{c}_{aa}^{\check{}} \end{bmatrix} \begin{bmatrix} \check{\Phi}_{a-}^1 \\ \check{\Phi}_a^1 \\ \check{\Phi}_a^1 \end{bmatrix} \\ &= \begin{bmatrix} \bar{e}_{aa-}^{\check{}} \bar{b}_{aa-}^{\check{}} & \bar{c}_{aa}^{\check{}} & \bar{e}_{aa}^{\check{}} \bar{b}_{aa}^{\check{}} \end{bmatrix} \begin{bmatrix} \check{\Phi}_{a-}^1 \\ \check{\Phi}_a^1 \\ \check{\Phi}_a^1 \end{bmatrix} \\ &= \begin{bmatrix} \bar{e}_{aa-}^{\check{}} & \bar{c}_{aa}^{\check{}} & \bar{e}_{aa}^{\check{}} \end{bmatrix} \begin{bmatrix} \alpha_a^- \\ \bar{\Phi}_a^1 \\ \alpha_a^+ \end{bmatrix} \end{aligned}$$

Since

$$\begin{bmatrix} \alpha_a^- \\ \bar{\Phi}_a^1 \\ \alpha_a^+ \end{bmatrix}$$

is an orthonormal basis for  $A_a^- \oplus \bar{V}_a^1 \oplus A_a^+$ , the matrix

$$\begin{bmatrix} \bar{e}_{aa-}^{\check{}} & \bar{c}_{aa}^{\check{}} & \bar{e}_{aa}^{\check{}} \end{bmatrix}$$

has orthonormal rows.

Again, from Corollary 9 we have

$$\bar{W}_a := (A_a^- \oplus \bar{V}_a^1 \oplus A_a^+) \ominus \bar{V}_a^0.$$

If  $\bar{W}_a \neq \{0\}$  it follows that the matrices  $\bar{d}_{aa}^-$  and  $\bar{d}_{aa'}^\sim$  may be found by completing the matrix  $\begin{bmatrix} \bar{e}_{aa-}^\sim & \bar{c}_{aa}^\sim & \bar{e}_{aa}^\sim \end{bmatrix}$  to an orthogonal square matrix, i.e., by determining matrices  $\bar{f}_{aa-}^\sim$ ,  $\bar{d}_{aa}^-$ , and  $\bar{f}_{aa}^\sim$  such that

$$\begin{bmatrix} \bar{e}_{aa-}^\sim & \bar{c}_{aa}^\sim & \bar{e}_{aa}^\sim \\ \bar{f}_{aa-}^\sim & \bar{d}_{aa}^- & \bar{f}_{aa}^\sim \end{bmatrix}$$

is an orthogonal matrix. Thus we have

$$\begin{bmatrix} \bar{\Phi}_a^0 \\ \bar{\Psi}_a \end{bmatrix} = \begin{bmatrix} \bar{e}_{aa-}^\sim & \bar{c}_{aa}^\sim & \bar{e}_{aa}^\sim \\ \bar{f}_{aa-}^\sim & \bar{d}_{aa}^- & \bar{f}_{aa}^\sim \end{bmatrix} \begin{bmatrix} \alpha_a^- \\ \bar{\Phi}_a^1 \\ \alpha_a^+ \end{bmatrix} = \begin{bmatrix} \bar{c}_{aa-}^\sim & \bar{c}_{aa}^\sim & \bar{c}_{aa}^\sim \\ \bar{d}_{aa-}^\sim & \bar{d}_{aa}^- & \bar{d}_{aa}^\sim \end{bmatrix} \begin{bmatrix} \bar{\Phi}_{a-}^1 \\ \bar{\Phi}_a^1 \\ \bar{\Phi}_a^1 \end{bmatrix},$$

where  $\bar{d}_{aa'}^\sim = \bar{f}_{aa'}^\sim \bar{b}_{aa'}^\sim$  for  $a' = a_-, a$ . If  $\bar{W}_a = \{0\}$ , then  $\bar{\Psi}_a$  is empty.

For each knot  $a$ , define

$$c_a := \begin{bmatrix} \bar{c}_{aa-}^\sim & \bar{c}_{aa}^\sim & \bar{c}_{aa}^\sim \\ 0 & 0 & \bar{c}_{aa}^\sim \end{bmatrix}.$$

Similarly, define

$$d_a := \begin{bmatrix} \bar{d}_{aa-}^\sim & \bar{d}_{aa}^- & \bar{d}_{aa}^\sim \\ 0 & 0 & \bar{d}_{aa}^\sim \end{bmatrix}.$$

By equations (42) and (43) we have that

$$\begin{bmatrix} \bar{\Phi}_a^0 \\ \bar{\Phi}_a^0 \end{bmatrix} = c_a \begin{bmatrix} \bar{\Phi}_{a-}^1 \\ \bar{\Phi}_a^1 \\ \bar{\Phi}_a^1 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} \bar{\Psi}_a \\ \bar{\Psi}_a \end{bmatrix} = d_a \begin{bmatrix} \bar{\Phi}_{a-}^1 \\ \bar{\Phi}_a^1 \\ \bar{\Phi}_a^1 \end{bmatrix}.$$

It follows from Corollary 9 that

$$V_a^1 = A_{a-}^+ \oplus V_a^0 \oplus W_a \oplus A_{a+}^-.$$

Since

$$\begin{bmatrix} \alpha_{a-}^+ \\ \bar{\Phi}_{a-}^0 \\ \bar{\Phi}_a^0 \\ \bar{\Phi}_a^0 \\ \bar{\Psi}_{a-} \\ \bar{\Psi}_a \\ \bar{\Psi}_a \\ \alpha_{a+}^- \end{bmatrix} = \begin{bmatrix} \bar{b}_{a-a-}^\sim & 0 & 0 \\ \bar{c}_{a-a-}^\sim & 0 & 0 \\ \bar{c}_{aa-}^\sim & \bar{c}_{aa}^\sim & \bar{c}_{aa}^\sim \\ 0 & 0 & \bar{c}_{aa}^\sim \\ \bar{d}_{a-a-}^\sim & 0 & 0 \\ \bar{d}_{aa-}^\sim & \bar{d}_{aa}^- & \bar{d}_{aa}^\sim \\ 0 & 0 & \bar{d}_{aa}^\sim \\ 0 & 0 & \bar{b}_{a+a}^\sim \end{bmatrix} \begin{bmatrix} \bar{\Phi}_{a-}^1 \\ \bar{\Phi}_a^1 \\ \bar{\Phi}_a^1 \end{bmatrix}$$

it follows that the matrix

$$\begin{bmatrix} \bar{b}_{a-a-}^\sim & 0 & 0 \\ \bar{c}_{a-a-}^\sim & 0 & 0 \\ \bar{c}_{aa-}^\sim & \bar{c}_{aa}^\sim & \bar{c}_{aa}^\sim \\ 0 & 0 & \bar{c}_{aa}^\sim \\ \bar{d}_{a-a-}^\sim & 0 & 0 \\ \bar{d}_{aa-}^\sim & \bar{d}_{aa}^- & \bar{d}_{aa}^\sim \\ 0 & 0 & \bar{d}_{aa}^\sim \\ 0 & 0 & \bar{b}_{a+a}^\sim \end{bmatrix} = \begin{bmatrix} \bar{b}_{a-a-}^\sim & 0 & 0 \\ \bar{c}_{a-a-}^\sim & 0 & 0 \\ & \bar{c}_a & \\ \bar{d}_{a-a-}^\sim & 0 & 0 \\ & \bar{d}_a & \\ 0 & 0 & \bar{b}_{a+a}^\sim \end{bmatrix}$$

is orthogonal.

We can generalize this orthogonal matrix as follows. Let  $a < b$  be knots so that  $[a, b]$  contains at least three knots. We define matrices  $c_{[a,b]}$  inductively:

$$c_{[a-, a+]} := \begin{bmatrix} \check{c}_{a- a-} & 0 & 0 \\ & c_a & \end{bmatrix}.$$

Suppose  $[a, b]$  contains more than three knots. Then

$$c_{[a,b]} := \begin{bmatrix} c_{[a,b-]} & 0 & 0 \\ 0 \dots c_{b-} & & \end{bmatrix}.$$

Note that the above notation indicates that the second row of the block array consists of zeros followed by  $c_{b-}$  at the end. For example

$$c_{[a-, a+]} = \begin{bmatrix} \check{c}_{a- a-} & 0 & 0 \\ \check{c}_{aa-} & \check{c}_{aa} & \check{c}_{aa} \\ 0 & 0 & c_{aa} \end{bmatrix}.$$

For four consecutive knots  $a < b < c < d$ ,

$$c_{[a,d]} = \begin{bmatrix} \check{c}_{aa} & 0 & 0 & 0 & 0 \\ \check{c}_{ba} & \check{c}_{bb} & \check{c}_{bb} & 0 & 0 \\ 0 & 0 & \check{c}_{bb} & 0 & 0 \\ 0 & 0 & c_{cb} & \check{c}_{cc} & \check{c}_{cc} \\ 0 & 0 & 0 & 0 & c_{cc} \end{bmatrix}.$$

Similarly we defined matrices  $d_{[a,b]}$  inductively.

$$d_{[a-, a+]} := \begin{bmatrix} \check{d}_{a- a-} & 0 & 0 \\ & d_a & \end{bmatrix}.$$

If  $[a, b]$  contains more than three knots, then

$$d_{[a,b]} := \begin{bmatrix} d_{[a,b-]} & 0 & 0 \\ 0 \dots d_{b-} & & \end{bmatrix}.$$

Next, we use these matrices to define

$$M_{[a,b]} := \begin{bmatrix} \check{b}_{aa} & 0 \dots 0 \\ c_{[a,b]} \\ d_{[a,b]} \\ 0 \dots 0 \check{b}_{bb-} \end{bmatrix}$$

It follows that

$$\begin{bmatrix} \alpha_a^+ \\ \check{\Phi}_a^0 \\ \check{\Phi}_{a+}^0 \\ \check{\Phi}_{a+}^0 \\ \vdots \\ \check{\Phi}_{b-}^0 \\ \check{\Psi}_a \\ \check{\Psi}_{a+} \\ \check{\Psi}_{a+} \\ \vdots \\ \check{\Psi}_{b-} \\ \alpha_b^- \end{bmatrix} = M_{[a,b]} \begin{bmatrix} \check{\Phi}_a^1 \\ \check{\Phi}_{a+}^1 \\ \check{\Phi}_{a+}^1 \\ \vdots \\ \check{\Phi}_{b-}^1 \end{bmatrix},$$

and thus  $M_{[a,b]}$  is an orthogonal matrix.

We now use the results of Section 2.1 to decompose the matrix

$$\begin{bmatrix} \check{d}_{aa-} & \check{d}_{aa} & \check{d}_{aa} \end{bmatrix}$$

when it is defined.

Using the fact that  $\hat{W}_a = (I - P_{\check{V}_a^0})\check{V}_a^1$  (see Lemma 14), we obtain that

$$\begin{bmatrix} -(c_{aa}^-)^T(\check{c}_{aa-}^-) & (I - (c_{aa}^-)^T c_{aa}^-) & -(c_{aa}^-)^T(\check{c}_{aa}^-) \end{bmatrix} \begin{bmatrix} \check{\Phi}_{a-}^1 \\ \check{\Phi}_a^1 \\ \check{\Phi}_a^1 \end{bmatrix}$$

is a basis for  $\hat{W}_a$ . Let  $\hat{g}_a$  be a matrix with orthonormal rows with the same row span as

$$\begin{bmatrix} -(c_{aa}^-)^T(\check{c}_{aa-}^-) & (I - (c_{aa}^-)^T c_{aa}^-) & -(c_{aa}^-)^T(\check{c}_{aa}^-) \end{bmatrix}.$$

Write

$$\hat{g}_a = [\check{g}_{aa-} \quad \check{g}_{aa} \quad \check{g}_{aa}].$$

Then

$$[\check{g}_{aa-} \quad \check{g}_{aa} \quad \check{g}_{aa}] \begin{bmatrix} \check{\Phi}_{a-}^1 \\ \check{\Phi}_a^1 \\ \check{\Phi}_a^1 \end{bmatrix}$$

is an orthonormal basis for  $\hat{W}_a$ .

Also, from Lemma 16, we have that  $\tilde{W}_a = (I - P_{\check{V}_a^0 \oplus \hat{W}_a})(A_a^- \oplus A_a^+)$ . This implies that

$$\begin{bmatrix} \check{b}_{aa-}^- & 0 & 0 \\ 0 & 0 & \check{b}_{aa}^- \end{bmatrix} \left( I - \begin{bmatrix} \check{c}_{aa-}^- & \check{c}_{aa}^- & \check{c}_{aa}^- \\ \check{g}_{aa-}^- & \check{g}_{aa}^- & \check{g}_{aa}^- \end{bmatrix}^T \begin{bmatrix} \check{c}_{aa-}^- & \check{c}_{aa}^- & \check{c}_{aa}^- \\ \check{g}_{aa-}^- & \check{g}_{aa}^- & \check{g}_{aa}^- \end{bmatrix} \right) \begin{bmatrix} \check{\Phi}_{a-}^1 \\ \check{\Phi}_a^1 \\ \check{\Phi}_a^1 \end{bmatrix}$$

is a basis for  $\tilde{W}_a$ . Let  $\tilde{g}_a$  be a matrix with orthonormal rows and with the same row space as

$$\begin{bmatrix} \check{b}_{aa-}^- & 0 & 0 \\ 0 & 0 & \check{b}_{aa}^- \end{bmatrix} \left( I - \begin{bmatrix} \check{c}_{aa-}^- & \check{c}_{aa}^- & \check{c}_{aa}^- \\ \check{g}_{aa-}^- & \check{g}_{aa}^- & \check{g}_{aa}^- \end{bmatrix}^T \begin{bmatrix} \check{c}_{aa-}^- & \check{c}_{aa}^- & \check{c}_{aa}^- \\ \check{g}_{aa-}^- & \check{g}_{aa}^- & \check{g}_{aa}^- \end{bmatrix} \right)$$

Write

$$\tilde{g}_a = [\check{g}_{aa-}^- \quad \check{g}_{aa}^- \quad \check{g}_{aa}^-].$$

Then

$$\begin{bmatrix} \tilde{g}_{aa-} & \tilde{g}_{aa} & \tilde{g}_{aa} \end{bmatrix} \begin{bmatrix} \check{\Phi}_{a-}^1 \\ \check{\Phi}_a^1 \\ \check{\Phi}_a^1 \end{bmatrix}$$

is an orthonormal basis for  $\tilde{W}_a$ .

Thus we have that

$$\begin{bmatrix} d_{aa-}^\vee & d_{aa}^\vee & d_{aa}^\vee \end{bmatrix} = \begin{bmatrix} \tilde{g}_{aa-}^\vee & \tilde{g}_{aa}^\vee & \tilde{g}_{aa}^\vee \\ \tilde{g}_{aa-}^\vee & \tilde{g}_{aa}^\vee & \tilde{g}_{aa}^\vee \end{bmatrix}.$$

**2.4. Arbitrary polynomial reproduction: Wavelet construction.** For a knot sequence  $\mathbf{a}$  and  $n = 1, 2, \dots$ , let  $V^0 := S(\Omega_{\mathbf{a}}^n)$  and  $V^1 := S(\Omega_{\mathbf{a}}^{n+3})$  where  $\Omega_{\mathbf{a}}^n$  is given in Section 1.3. By Theorem 7 there is a basis  $\Psi_{\mathbf{a}}^n$  centered on  $\mathbf{a}$  such that  $V^1 = V^0 \oplus W$  where  $W = S(\Psi_{\mathbf{a}}^n)$ . We next use the algorithm from section 2.2 to construct such a  $\Psi_{\mathbf{a}}^n$ .

Recall that

$$r^n = (I - P_{\{\tilde{\phi}^2, \dots, \tilde{\phi}^n, z^n\}})r = (I - P_{\{z^n\}})r_n$$

and

$$l^n = (I - P_{\{\tilde{\phi}^2, \dots, \tilde{\phi}^n, z^n\}})l = (I - P_{\{z^n\}})l_n.$$

For  $n = 1, 2, \dots$ , let

$$\Lambda^n := \text{span}\{\tilde{\phi}^2, \dots, \tilde{\phi}^n, z^n\}.$$

Note that  $\Lambda^1 = \text{span}\{z^1\}$ .

From the definitions of  $\tilde{\phi}^n$  and  $z^n$  it follows that

$$\Lambda^{n+3} = \Lambda^n \oplus \text{span}\{\tilde{\phi}^{n+2}, z_{\perp}^n, z^{n+3}\} = \Lambda^n \oplus \Delta^n,$$

where

$$z_{\perp}^n = \frac{\tilde{\phi}^{n+1}}{\|\tilde{\phi}^{n+1}\|^2} - \alpha_n \frac{\tilde{\phi}^{n+3}}{\|\tilde{\phi}^{n+3}\|^2}$$

and

$$\Delta^n := \text{span}\{\tilde{\phi}^{n+2}, z_{\perp}^n, z^{n+3}\}.$$

Clearly,  $z_{\perp}^n$  and  $z^n$  are orthogonal and span the same space as  $\tilde{\phi}^{n+1}$  and  $\tilde{\phi}^{n+3}$ .

Now

$$\bar{V}_a^0 = \text{span } \bar{\Omega}_a^n = \text{span}\{\bar{\omega}_a^n\}$$

where  $\bar{\omega}_a^n = r^n \circ \sigma_{a-} + l^n \circ \sigma_a$ .

Similarly

$$\bar{V}_a^1 = \text{span } \bar{\Omega}_a^{n+3} = \text{span}\{\bar{\omega}_a^{n+3}\}$$

where  $\bar{\omega}_a^{n+3} = r^{n+3} \circ \sigma_{a-} + l^{n+3} \circ \sigma_a$ .

It follows that  $\bar{k}_a^0 = \bar{k}_a^1 = 1$  for every  $a$ . Furthermore it is easy to see that for every  $a$ ,  $Y_a = Y_a^- = Y_a^+ = \{0\}$ , and so  $m_a = m_a^- = m_a^+ = 0$ . Lemmas 14 and 16 imply that  $\dim \tilde{W}_a^n = \dim \tilde{W}_a^{n+3} = 1$ . The superscript denotes the dependence on  $n$  in the construction.

From the definitions of  $r^n$  and  $l^n$ , and the symmetries of the polynomials  $\tilde{\phi}^n$ , it can be shown that  $l^n$  is the reflection of  $r^n$  with respect to the line  $x = 1/2$ . From this we can see that

$$\langle \bar{\omega}_a^n, \bar{\omega}_a^n \rangle = (a_+ - a_-) \langle r^n, r^n \rangle$$

$$\langle \bar{\omega}_a^n, \bar{\omega}_a^{n+3} \rangle = (a_+ - a_-) \langle r^n, r^{n+3} \rangle$$

Since  $\Lambda^{n+3} = \Lambda^n \oplus \Delta^n$  and  $r^n = r - P_{\Lambda^n} r$ , we get

$$(46) \quad r^n - r^{n+3} = P_{\Delta^n} r.$$

Now,  $r^{n+3} \perp \Delta^n$ . From this it follows that

$$\langle r^n, r^{n+3} \rangle = \langle r^{n+3}, r^{n+3} \rangle.$$

We now begin the construction of the wavelet constructions. Since  $\hat{W}_a^n$  is one dimensional equation (39) shows it is spanned by the single function  $\hat{w}_a^n := (I - P_{\bar{w}_a^n}) \bar{\omega}_a^{n+3}$  which is,

$$\hat{w}_a^n = (r^{n+3} - \frac{\langle r^{n+3}, r^{n+3} \rangle}{\langle r^n, r^n \rangle} r^n) \circ \sigma_{a_-} + (l^{n+3} - \frac{\langle r^{n+3}, r^{n+3} \rangle}{\langle r^n, r^n \rangle} l^n) \circ \sigma_a.$$

Since  $T_a^n = (P_{\bar{V}_{a_-}^1} + P_{\bar{V}_a^1}) \bar{V}_a^0$  it is one dimensional and by Lemma 13 it follows that  $T_a^- = T_a^+ = \{0\}$ , and so  $U_a^n = T_a^n$  hence  $S_a^n$  is spanned by the single function

$$s_a^n = -(r^n - r^{n+3}) \circ \sigma_{a_-} + (l^n - l^{n+3}) \circ \sigma_a.$$

Equation (40) shows that  $\tilde{W}_a^n$  is spanned by  $(I - P_{\bar{V}_a^0 \oplus \bar{W}_a^n}) s_a^n$ . A straightforward computation shows that this function is a scalar multiple of

$$\tilde{w}_a^n := (r^n - r^{n+3}) \circ \sigma_{a_-} - c_a (l^n - l^{n+3}) \circ \sigma_a.$$

where

$$c_a = \frac{a - a_-}{a_+ - a_-}.$$

We now construct the short wavelets. Since  $\check{k}_a^0 = n$ ,  $\check{k}_a^1 = n + 3$  and from above  $\bar{k}_a^0 = \bar{k}_a^1 = 1$  it follows from Theorem 11 that  $\dim \check{W}_a^n = 1$ . Also from the formula for  $T_a^n$  above we find

$$\begin{aligned} A_a^- &= \text{span} \{ (r^n - r^{n+3}) \circ \sigma_{a_-} \} \\ A_a^+ &= \text{span} \{ (l^n - l^{n+3}) \circ \sigma_a \} \end{aligned}$$

With the above results, using equation (41) and another straightforward computation we see that  $\check{W}_a^n$  is spanned by

$$\check{w}_a^n := \langle z^{n+3}, r \rangle \tilde{\phi}^{n+2} \circ \sigma_a - \langle \tilde{\phi}^{n+2}, r \rangle z^{n+3} \circ \sigma_a.$$

In summary the wavelets constructed are,

$$(47) \quad \begin{aligned} \hat{w}_a^n &= (r^{n+3} - \frac{\langle r^{n+3}, r^{n+3} \rangle}{\langle r^n, r^n \rangle} r^n) \circ \sigma_{a_-} + (l^{n+3} - \frac{\langle r^{n+3}, r^{n+3} \rangle}{\langle r^n, r^n \rangle} l^n) \circ \sigma_a \\ \tilde{w}_a^n &= (r^n - r^{n+3}) \circ \sigma_{a_-} - c_a (l^n - l^{n+3}) \circ \sigma_a \\ \check{w}_a^n &= \langle z^{n+3}, r \rangle \tilde{\phi}^{n+2} \circ \sigma_a - \langle \tilde{\phi}^{n+2}, r \rangle z^{n+3} \circ \sigma_a. \end{aligned}$$

### 3. $\tau$ -WAVELETS

**3.1. Nested Knot Sequences Determined by  $\tau$ .** Let  $\tau$  denote the "golden ratio"  $\frac{1}{2}(1 + \sqrt{5})$  which satisfies the quadratic relation  $\tau^2 = 1 + \tau$ . A non-negative number  $x$  is a  **$\tau$ -rational number** if it can be represented in the form  $x = \sum_{k=m}^n \varepsilon_k \tau^k$  where  $m \leq n$  are integers and each  $\varepsilon_k \in \{0, 1\}$ . Furthermore,  $x$  is a  **$\tau$ -integer** if it has such a representation with  $m = 0$ . By the above quadratic relation, this representation is unique if one further requires that  $\varepsilon_k \varepsilon_{k+1} = 0$  whenever  $m \leq k \leq n-1$ . For example, the first few non-negative  $\tau$ -integers are  $0, 1, \tau, \tau^2, \tau^2 + 1, \tau^3, \dots$ . We denote the set of non-negative  $\tau$ -integers by  $\mathbf{Z}_\tau^+$ . The unique representation of positive  $\tau$ -integers implies that  $\mathbf{Z}_\tau^+ \setminus \{0\}$  can be partitioned into  $\tau \mathbf{Z}_\tau^+ \setminus \{0\}$  and  $1 + \tau^2 \mathbf{Z}_\tau^+$ . Further,  $\tau \mathbf{Z}_\tau^+ \setminus \{0\}$  can be partitioned into  $\tau + \tau^2 \mathbf{Z}_\tau^+$  and  $\tau^2 + \tau^3 \mathbf{Z}_\tau^+$ . Also, the difference between consecutive  $\tau$ -integers is either 1 or  $\frac{1}{\tau}$ . More specifically, considering  $\mathbf{Z}_\tau^+$  as a knot sequence  $\mathbf{a}$ , if  $a \in 1 + \tau^2 \mathbf{Z}_\tau^+$ , then  $a - a_- = 1$  and  $a_+ - a = \frac{1}{\tau}$ ; while if  $a \in \tau + \tau^2 \mathbf{Z}_\tau^+$ , then  $a - a_- = \frac{1}{\tau}$  and  $a_+ - a = 1$ ; and finally if  $a \in \tau^2 + \tau^3 \mathbf{Z}_\tau^+$ , then  $a - a_- = 1$  and  $a_+ - a = 1$ . Letting  $L$  denote the "long" difference of 1 and letting  $S$  denote the "short" difference of  $\frac{1}{\tau}$ , we denote a non-zero  $a \in \mathbf{a}$  as  $LS$ , respectively  $SL, LL$ , if  $a \in 1 + \tau^2 \mathbf{Z}_\tau^+$ , respectively  $a \in \tau + \tau^2 \mathbf{Z}_\tau^+$ ,  $a \in \tau^2 + \tau^3 \mathbf{Z}_\tau^+$ . (Note: The sequence of successive differences of elements of  $\mathbf{Z}_\tau^+$  forms an infinite word,

$$f = LSLLSLSLLSLLSLSLLSLSL \dots,$$

with alphabet  $\{L, S\}$  which is invariant under the substitution  $L \mapsto LS, S \mapsto L$ .  $f$  is called the *Fibonacci word*; see [19].)

For  $k \in \mathbf{Z}$ , let  $\mathbf{a}^k$  denote the knot sequence  $(\frac{1}{\tau})^k \mathbf{Z}_\tau^+$ . The above unique representation of  $\tau$ -integers shows that  $\mathbf{a}^k \subset \mathbf{a}^{k+1}$  for  $k \in \mathbf{Z}$ . Also,  $\bigcup_{k \in \mathbf{Z}} \mathbf{a}^k$  is the set of non-negative  $\tau$ -rational numbers which can be shown to be dense in  $\mathbf{R}^+$ . For an integer  $k$  and  $a \in \mathbf{a}^k$ , let  $a^k$  (resp.  $a_+^k$ ) denote the predecessor (resp. successor) of  $a$  relative to  $\mathbf{a}^k$ . At level  $k$ , each interval  $[a, a_+^k]$  has length either  $\frac{1}{\tau^k}$  or  $\frac{1}{\tau^{k+1}}$ . Such an interval will be called long at level  $k$  if it has length  $\frac{1}{\tau^k}$  and short at level  $k$  if it has length  $\frac{1}{\tau^{k+1}}$ . The refinement from level  $k$  to level  $k+1$  proceeds as follows. Each long interval at level  $k$  is split into two subintervals  $[a, a_+^{k+1}]$  and  $[a_+^{k+1}, a_+^k]$ , where

$$(48) \quad a_+^{k+1} = (1 - \frac{1}{\tau})a + \frac{1}{\tau}a_+^k.$$

It follows that the left subinterval  $[a, a_+^{k+1}]$  is long at level  $k+1$ , and the right subinterval  $[a_+^{k+1}, a_+^k]$  is short at level  $k+1$ . Each short interval at level  $k$  is not subdivided and becomes long at level  $k+1$ ; i.e. if  $a_+^k - a = \frac{1}{\tau^{k+1}}$  then  $a_+^k = a_+^{k+1}$ .

**3.2.  $\tau$ -Wavelets of Haar.** We now show that the  $\tau$ -wavelets of Haar, constructed in [14], can be considered as a special case of the general wavelet construction outlined in Section 2.2. Let  $\phi_1 := \chi_{[0,1]}$ ,  $\phi_2 := \tau^{\frac{1}{2}} \chi_{[0, \frac{1}{\tau}]}$ , and  $\Phi = \{\phi_1(\cdot - b) \mid b \in \tau \mathbf{Z}_\tau^+\} \cup \{\phi_2(\cdot - c) \mid c \in 1 + \tau^2 \mathbf{Z}_\tau^+\}$ . Then  $\Phi$  is an orthonormal basis centered on  $\mathbf{Z}_\tau^+$  that is generated by appropriate  $\tau$ -integer translations of  $\phi_1$  and  $\phi_2$  and  $S(\Phi)$  is the space of piecewise constant functions in  $L^2(\mathbf{R}^+)$  with breakpoints in  $\mathbf{Z}_\tau^+$ . For  $k \in \mathbf{Z}$ , let  $\Phi^k := \{\tau^{\frac{k}{2}} f(\tau^k \cdot) \mid f \in \Phi\} = \{\tau^{\frac{k}{2}} \phi_1(\tau^k \cdot - b) \mid b \in \tau \mathbf{Z}_\tau^+\} \cup \{\tau^{\frac{k}{2}} \phi_2(\tau^k \cdot - c) \mid$

$c \in 1 + \tau^2 \mathbf{Z}_\tau^+$ ). Then  $\Phi^k$  is an orthonormal basis centered on  $\mathbf{a}^k$  and  $S(\Phi^k)$  is the space of piecewise constant functions in  $L^2(\mathbf{R}^+)$  with breakpoints in  $\mathbf{a}^k$ .

Since  $\mathbf{a}^k \subset \mathbf{a}^{k+1}$  for  $k \in \mathbf{Z}$ , it follows that  $S(\Phi^k) \subset S(\Phi^{k+1})$  for  $k \in \mathbf{Z}$ . In particular,  $\Phi^0$  and  $\Phi^1$  are orthonormal bases centered on  $\mathbf{Z}_\tau^+$  and  $S(\Phi^0) \subset S(\Phi^1)$ . Theorem 7 thus applies to this situation. We construct  $\Psi$  as outlined in Section 2.2.

For  $a > 0 \in \mathbf{Z}_\tau^+$ , it is easy to see that  $\bar{k}_a^0 = \bar{k}_a^1 = m_a = m_a^\pm = 0$ . Also,  $\check{k}_a^0 = 1$  for every  $a \in \mathbf{Z}_\tau^+$ . For  $a \in \mathbf{Z}_\tau^+$ , it follows from the manner in which  $\mathbf{a}^0$  is refined to  $\mathbf{a}^1$  that

$$\check{k}_a^1 = \begin{cases} 1 & \text{if } a \in 1 + \tau^2 \mathbf{Z}_\tau^+ \\ 2 & \text{if } a \in \tau \mathbf{Z}_\tau^+. \end{cases}$$

Thus Theorem 11 implies that  $\bar{W}_a = \{0\}$  for  $a > 0 \in \mathbf{Z}_\tau^+$ , and

$$\dim \check{W}_a = \begin{cases} 0 & \text{if } a \in 1 + \tau^2 \mathbf{Z}_\tau^+ \\ 1 & \text{if } a \in \tau \mathbf{Z}_\tau^+. \end{cases}$$

From equation (35), for  $a \in \tau \mathbf{Z}_\tau^+$  we have  $\check{W}_a = (I - P_{\check{V}_a^0}) \check{V}_a^1$ . Also, for  $a \in \tau \mathbf{Z}_\tau^+$ ,  $\check{V}_a^0$  is spanned by the function  $\chi_{[a, a_+]}$  and  $\check{V}_a^1$  is spanned by the functions  $\chi_{[a, a + \frac{1}{\tau}]}$  and  $\chi_{[a + \frac{1}{\tau}, a_+]}$ . An easy computation shows that an orthonormal basis for  $\check{W}_a$  consists of the single function  $\tau^{-\frac{1}{2}} \chi_{[a, a + \frac{1}{\tau}]} - \tau^{\frac{1}{2}} \chi_{[a + \frac{1}{\tau}, a_+]}$ .

If we define

$$\psi := \tau^{-\frac{1}{2}} \chi_{[0, \frac{1}{\tau}]} - \tau^{\frac{1}{2}} \chi_{[\frac{1}{\tau}, 1]},$$

it follows that  $W = S(\Psi)$  where  $\Psi := \{\psi(\cdot - b) \mid b \in \tau \mathbf{Z}_\tau^+\}$ . Note that  $\Psi$  is an orthonormal basis centered on  $\mathbf{Z}_\tau^+$ , obtained by appropriate  $\tau$ -integer translations of  $\psi$ .

Letting  $\Psi^k := \{\tau^{\frac{k}{2}} \psi(\tau^k \cdot -b) \mid b \in \tau \mathbf{Z}_\tau^+\}$ , for  $k \in \mathbf{Z}$ , it is easy to check that  $\Psi^k$  is a basis centered on  $\mathbf{a}^k$  and that  $S(\Phi^k) \oplus S(\Psi^k) = S(\Phi^{k+1})$ . Since  $\bigcap_{k \in \mathbf{Z}} S(\Phi^k) = \{0\}$  and  $\bigcup_{k \in \mathbf{Z}} S(\Phi^k)$  is dense in  $L^2(\mathbf{R}^+)$ , it follows that  $L^2(\mathbf{R}^+) = \bigoplus_{k \in \mathbf{Z}} S(\Psi^k)$ , i.e.  $\{\tau^{\frac{k}{2}} \psi(\tau^k \cdot -b) \mid k \in \mathbf{Z}, b \in \tau \mathbf{Z}_\tau^+\}$  is an orthonormal basis of  $L^2(\mathbf{R}^+)$ . These functions are called the  $\tau$ -wavelets of Haar in [14].

### 3.3. Continuous, Piecewise Quadratic $\tau$ -Wavelets. Let

$$(49) \quad \Phi^k := \Omega_{\mathbf{a}^k, \boldsymbol{\theta}}, \quad k \in \mathbf{Z}$$

denote the continuous, piecewise quadratic, orthonormal basis centered on  $\mathbf{a}^k$  as described in Section 1.4 with constant parameter sequence  $\boldsymbol{\theta} = (\theta_a)_{a \in \mathbf{a}^k}$  where  $\theta_a = 1/\tau$  for  $a \in \mathbf{a}^k$  (here we assume that the components of  $\Phi^k$  have been normalized). It follows from (48) that  $(\mathbf{a}^k, \boldsymbol{\theta})$  satisfy the hypotheses of Lemma 6. Since, as previously discussed  $\bigcup_k \mathbf{a}^k$  is dense in  $J = \mathbf{R}_+$  and since  $\boldsymbol{\alpha} = \bigcap_k \mathbf{a}^k \cup \mathbf{b}(\mathbf{a}^k, \boldsymbol{\theta}^k) = \{0\}$ , Lemma 6 implies that the spaces  $V^k := S(\Phi^k)$  form a **multiresolution analysis of  $L^2(\mathbf{R}_+)$** , that is,

- (1)  $V^k \subset V^{k+1}$ ,  $(k \in \mathbf{Z})$ ,
- (2)  $\bigcup_k V^k$  is dense in  $L^2(\mathbf{R}_+)$ ,
- (3)  $\bigcap_k V^k = \{0\}$ .

As in the  $\tau$ -wavelets of Haar construction, the basis  $\Phi := \Phi^0$  centered on the knot sequence  $\mathbf{a}^0 = \mathbf{Z}_\tau^+$  for  $V_0$  can also be generated by the  $\tau$ -integer translations of a small number of functions in  $\Phi$ . Suppose  $a$  and  $a'$  in  $\mathbf{Z}_\tau^+ \setminus \{0\}$  have the same

classification (say  $LS$ ), then from the construction of  $\Phi$  it follows that  $\Phi_a$  is a translate of  $\Phi_{a'}$ . (Note: Here  $\Phi_a$  stands for the more cumbersome, yet precise,  $\Phi_{a,\mathbf{a}^0}^0$ ; see section 1.) Specifically, we have, for  $b \in \mathbf{Z}_\tau^+$ ,

$$(50) \quad \begin{aligned} \Phi_{1+\tau^2 b}(\cdot) &= \Phi_1(\cdot - \tau^2 b), \\ \Phi_{\tau+\tau^2 b}(\cdot) &= \Phi_\tau(\cdot - \tau^2 b), \\ \Phi_{\tau^2+\tau^3 b}(\cdot) &= \Phi_{\tau^2}(\cdot - \tau^3 b). \end{aligned}$$

Thus every function in  $\Phi$ , with the exception of those in  $\Phi_0$ , is a  $\tau$ -integer translate of a function from  $\Phi_1$ ,  $\Phi_\tau$ , or  $\Phi_{\tau^2}$ . For any positive  $a \in \mathbf{Z}_\tau^+$ , let

$$\beta(a) := \begin{cases} 1 & \text{if } a \in 1 + \tau^2 \mathbf{Z}_\tau^+, \\ \tau & \text{if } a \in \tau + \tau^2 \mathbf{Z}_\tau^+, \\ \tau^2 & \text{if } a \in \tau^2 + \tau^3 \mathbf{Z}_\tau^+, \end{cases}$$

and  $\mu(a) := a - \beta(a)$ . Then  $a = \beta(a) + \mu(a)$  where  $\beta(a) \in \{1, \tau, \tau^2\}$  and  $\mu(a) \in \tau^2 \mathbf{Z}_\tau^+$  if  $\beta(a) = 1$  or  $\tau$  and  $\mu(a) \in \tau^3 \mathbf{Z}_\tau^+$  if  $\beta(a) = \tau^2$ . Thus, we can write (50) in the more compact form

$$(51) \quad \Phi_a = \Phi_{\beta(a)}(\cdot - \mu(a)) \quad (a \in \mathbf{Z}_\tau^+ \setminus \{0\}).$$

For  $k \in \mathbf{Z}$  and  $a \in \mathbf{Z}_\tau^+$ , let

$$\Phi_{k,a} = \tau^{k/2} \Phi_a(\tau^k \cdot).$$

We remark that with this definition,  $\Phi_{k,a} = \Phi_{a/\tau^k, \mathbf{a}^k}^k$ , and we apologize for the confusing notation.

Note that  $V^0 \subset V^1$  and both  $\Phi^0$  and  $\Phi^1$  can be regarded as orthonormal bases centered on the knot sequence  $\mathbf{a}^0 = \mathbf{Z}_\tau^+$ . For  $a \in \mathbf{a}^0$ ,  $\Phi_a^1$  will denote the more precise  $\Phi_{a,\mathbf{a}^0}^1$ .

It follows that we have the following ‘polyphase’ like representation:

$$\Phi_0^1 = \begin{bmatrix} \Phi_{1,0} \\ \Phi_{1,1} \end{bmatrix}, \Phi_1^1 = \Phi_{1,\tau}, \Phi_\tau^1 = \begin{bmatrix} \Phi_{1,\tau^2} \\ \Phi_{1,1+\tau^2} \end{bmatrix}, \text{ and } \Phi_{\tau^2}^1 = \begin{bmatrix} \Phi_{1,\tau^3} \\ \Phi_{1,1+\tau^3} \end{bmatrix}.$$

For example, Figure 1 shows the three functions in  $\Phi_\tau$ .





FIGURE 1.  $\Phi_\tau$ .

Figure 2 shows the six functions in  $\Phi_\tau^1$ .

FIGURE 2.  $\Phi_\tau^1$ .

For  $a \in \mathbf{Z}_\tau^+$  and  $a > \tau^2$ ,

$$\Phi_a^1 = \Phi_{\beta(a)}^1(\cdot - \mu(a)).$$

The coefficient matrices from section 2.3 can now be computed. For  $a \in \mathbf{Z}_\tau^+$  and  $a' \in \{a_-, a\}$ ,  $c_{aa'} = \langle \Phi_a, \Phi_{a'}^1 \rangle$ . In particular we have

$$\Phi_0 = c_{00}\Phi_0^1, \text{ and } \Phi_a = c_{aa-}\Phi_{a-}^1 + c_{aa}\Phi_a^1, \text{ for } a > 0.$$

For any  $k \in \mathbf{Z}$ ,  $V^k \subset V^{k+1}$  and both  $\Phi^k$  and  $\Phi^{k+1}$  can be regarded as orthonormal bases centered on the knot sequence  $\mathbf{a}^k$ . For  $a \in \mathbf{Z}_\tau^+$  it follows that  $\Phi_{a/\tau^k, \mathbf{a}^k}^{k+1} = \tau^{k/2}\Phi_a^1(\tau^k \cdot)$ . We have, for  $b \in \mathbf{a}^k$ ,

$$\Phi_{0, \mathbf{a}^k}^k = c_{00}\Phi_{0, \mathbf{a}^k}^{k+1}, \text{ and } \Phi_{b, \mathbf{a}^k}^k = c_{(\tau^k b)(\tau^k b)-}\Phi_{b-, \mathbf{a}^k}^{k+1} + c_{(\tau^k b)(\tau^k b)}\Phi_{b, \mathbf{a}^k}^{k+1} \text{ for } b > 0.$$

(Note: In the second equation above there are two occurrences of the minus subscript. The first denotes the predecessor of  $\tau^k b$  in the knot sequence  $\mathbf{a}^0$ . The second denotes the predecessor of  $b$  in the knot sequence  $\mathbf{a}^k$ . See section 1.)

Recall that

$$\Phi_0^1 = \begin{bmatrix} \Phi_{1,0} \\ \Phi_{1,1} \end{bmatrix}.$$

Thus

$$c_{00} = \langle \Phi_0, \Phi_0^1 \rangle = [\langle \Phi_0, \Phi_{1,0} \rangle \quad \langle \Phi_0, \Phi_{1,1} \rangle].$$

This suggests defining, for  $a, a' \in \mathbf{Z}_\tau^+$ ,

$$C_{a,a'} := \langle \Phi_a, \Phi_{1,a'} \rangle.$$

We now have

$$c_{00} = [C_{0,0} \quad C_{0,1}] \text{ and } c_{10} = [C_{1,0} \quad C_{1,1}].$$

Similarly it follows that  $c_{11} = C_{1,\tau}$ ,  $c_{\tau 1} = 0$ ,

$$c_{\tau\tau} = [C_{\tau,\tau^2} \quad C_{\tau,1+\tau^2}], c_{\tau^2\tau} = [C_{\tau^2,\tau^2} \quad C_{\tau^2,1+\tau^2}],$$

and

$$c_{\tau^2\tau^2} = [C_{\tau^2,\tau^3} \quad C_{\tau^2,1+\tau^3}].$$

For  $a > \tau^2$ ,  $c_{aa} = c_{\beta(a)\beta(a)}$  and  $c_{aa-} = c_{\beta(a)\beta(a)-}$ .

In Table 1 we provide the matrices  $C_{a,a'}$ , mentioned above, which were computed with the aid of *Mathematica*<sup>TM</sup>. Here we assume that  $\Phi_a$  is ordered as follows: of the three components, the second and third components are the “ $q$ ” and “ $z$ ”

components from  $\check{\Phi}_a$ , respectively, while the first component is the component of  $\Phi_a$  that does not vanish at  $a$ .

TABLE 1. Scaling Coefficients for  $\Phi$ 

$$\begin{aligned}
C_{0,0} &= \begin{pmatrix} \sqrt{-\frac{1}{2} + \frac{\sqrt{5}}{2}} & -\frac{1}{2\sqrt{16491+7375\sqrt{5}}} & \frac{121}{2\sqrt{5485+2441\sqrt{5}}} \\ 0 & \frac{1}{4}\sqrt{-818+370\sqrt{5}} & -\sqrt{\frac{55}{2(405+181\sqrt{5})}} \\ 0 & \frac{251}{\sqrt{76350+33982\sqrt{5}}} & \frac{59}{\sqrt{30410+13610\sqrt{5}}} \end{pmatrix} \\
C_{0,1} &= \begin{pmatrix} \frac{19\sqrt{\frac{2}{9765+4387\sqrt{5}}}}{\sqrt{\frac{2}{11}(-35+16\sqrt{5})}} & \frac{1}{2}\sqrt{\frac{11}{767+343\sqrt{5}}} & \frac{19}{2}\sqrt{\frac{5}{25615+11463\sqrt{5}}} \\ -\frac{1}{11}\sqrt{\frac{5922}{5}-\frac{2584}{\sqrt{5}}} & -\sqrt{-\frac{283}{8}+\frac{637}{8\sqrt{5}}} & \frac{5}{\sqrt{370+166\sqrt{5}}} \\ 0 & 0 & 0 \end{pmatrix} \\
C_{1,0} &= \begin{pmatrix} 0 & -\frac{1}{2}\sqrt{\frac{11}{1715+767\sqrt{5}}} & \frac{11}{\sqrt{2}(35+17\sqrt{5})} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
C_{1,1} &= \begin{pmatrix} -\sqrt{-\frac{2}{5}+\frac{1}{\sqrt{5}}} & \frac{1}{4}\sqrt{177-\frac{389}{\sqrt{5}}} & -\frac{1}{4}\sqrt{-89+41\sqrt{5}} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
C_{1,\tau} &= \begin{pmatrix} 3-\sqrt{5} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \\
C_{\tau,\tau^2} &= \begin{pmatrix} \frac{1}{2}\sqrt{\frac{1}{2}(13-3\sqrt{5})} & -\frac{1}{4}\sqrt{\frac{5}{36875+16491\sqrt{5}}} & \frac{121}{4\sqrt{2441+1097\sqrt{5}}} \\ 0 & \sqrt{-\frac{409}{8}+\frac{185\sqrt{5}}{8}} & -\sqrt{\frac{55}{2(405+181\sqrt{5})}} \\ 0 & \frac{251}{\sqrt{76350+33982\sqrt{5}}} & \frac{59}{\sqrt{30410+13610\sqrt{5}}} \end{pmatrix} \\
C_{\tau,\tau^2+1} &= \begin{pmatrix} \frac{1}{44}(-63+31\sqrt{5}) & \frac{1}{4}\sqrt{\frac{55}{1715+767\sqrt{5}}} & \frac{19}{4}\sqrt{\frac{5}{11463+5123\sqrt{5}}} \\ \sqrt{\frac{2}{11}(-35+16\sqrt{5})} & \frac{1}{4}(13-5\sqrt{5}) & \frac{5}{\sqrt{370+166\sqrt{5}}} \\ -\frac{1}{11}\sqrt{\frac{5922}{5}-\frac{2584}{\sqrt{5}}} & -\sqrt{-\frac{283}{8}+\frac{637}{8\sqrt{5}}} & \frac{1}{44}(-129+53\sqrt{5}) \end{pmatrix} \\
C_{\tau^2,\tau^2} &= \begin{pmatrix} 0 & -\frac{1}{2}\sqrt{\frac{11}{2(1030+461\sqrt{5})}} & \frac{1}{2}\sqrt{\frac{1}{310}(166-69\sqrt{5})} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
C_{\tau^2,\tau^2+1} &= \begin{pmatrix} -\sqrt{\frac{1}{310}+\frac{\sqrt{5}}{62}} & \frac{1}{2}\sqrt{\frac{1}{310}(625-192\sqrt{5})} & -\sqrt{-\frac{37}{248}+\frac{4\sqrt{5}}{31}} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\
C_{\tau^2,\tau^3} &= \begin{pmatrix} 2\sqrt{\frac{1}{31}(6-\sqrt{5})} & -\frac{1}{2}\sqrt{\frac{5}{2(93850+41971\sqrt{5})}} & \frac{121}{2\sqrt{12492+5554\sqrt{5}}} \\ 0 & \sqrt{-\frac{409}{8}+\frac{185\sqrt{5}}{8}} & -\sqrt{\frac{55}{2(405+181\sqrt{5})}} \\ 0 & \frac{251}{\sqrt{76350+33982\sqrt{5}}} & \frac{59}{\sqrt{30410+13610\sqrt{5}}} \end{pmatrix} \\
C_{\tau^2,\tau^3+1} &= \begin{pmatrix} \frac{19\sqrt{\frac{2}{22219+9991\sqrt{5}}}}{\sqrt{\frac{2}{11}(-35+16\sqrt{5})}} & \frac{1}{2}\sqrt{\frac{55}{2(4365+1952\sqrt{5})}} & \frac{19}{2}\sqrt{\frac{5}{58306+26096\sqrt{5}}} \\ -\frac{1}{11}\sqrt{\frac{5922}{5}-\frac{2584}{\sqrt{5}}} & -\sqrt{-\frac{283}{8}+\frac{637}{8\sqrt{5}}} & \frac{5}{\sqrt{370+166\sqrt{5}}} \\ 0 & 0 & 0 \end{pmatrix}
\end{aligned}$$

3.3.1. *Construction of  $\tau$  wavelets.* We now construct an orthogonal wavelet basis  $\Psi$ , centered on  $\mathbf{a}^0$ , for the continuous, piecewise quadratic scaling functions  $\Phi$  such that

$$S(\Phi^0) \oplus S(\Psi) = S(\Phi^1)$$

using the method from section 2.2. As mentioned above this provides an example where the spaces  $T_a^\pm$  contain nonzero functions.

As in Section 2.2, we begin with determining the dimensions  $\bar{k}_a^\varepsilon$ ,  $m_a$ , and  $m_a^\pm$ .

If  $a > 0$  then  $\bar{k}_a^0 = \bar{k}_a^1 = 1$  and  $m_a = 0$ . Also, for  $a > 0$ ,

$$(52) \quad m_a^+ = \begin{cases} 1 & \text{if } a \in 1 + \tau^2 \mathbf{Z}_\tau^+ \\ 0 & \text{otherwise} \end{cases}$$

and

$$(53) \quad m_a^- = \begin{cases} 1 & \text{if } a \in \tau + \tau^2 \mathbf{Z}_\tau^+ \\ 0 & \text{otherwise.} \end{cases}$$

From Theorem 11 we get

$$\dim \bar{W}_a = \begin{cases} 1 & \text{if either } a \in 1 + \tau^2 \mathbf{Z}_\tau^+ \text{ or } a \in \tau + \tau^2 \mathbf{Z}_\tau^+ \\ 2 & \text{if } a \in \tau^2 + \tau^3 \mathbf{Z}_\tau^+. \end{cases}$$

It follows that for  $a \in 1 + \tau^2 \mathbf{Z}_\tau^+$  or  $a \in \tau + \tau^2 \mathbf{Z}_\tau^+$  that  $\bar{W}_a = \hat{W}_a$  which is spanned by a single function. For  $a \in \tau^2 + \tau^3 \mathbf{Z}_\tau^+$ , a basis for  $\bar{W}_a$  consists of two functions, one forming a basis for  $\hat{W}_a$  and the other a basis for  $\tilde{W}_a$ .

In all cases, Lemma 14 implies that for  $a > 0$ ,  $\dim \hat{W}_a = 1$ . To find a basis for  $\hat{W}_a$  we begin by constructing bases for  $\hat{W}_1, \hat{W}_\tau$ , and  $\hat{W}_{\tau^2}$ . We use the notation introduced in Section 1.4.

For  $a > 0$  an orthonormal basis of  $\bar{V}_a^0$  consists of the single function  $\bar{\phi}_a^0$  which is a suitable scalar multiple of  $r^{1/\tau} \circ \sigma_{a-} + l^{1/\tau} \circ \sigma_a$ . It follows that an orthonormal basis of  $\bar{V}_1^1$  consists of a single function  $\bar{\phi}_1^1$  which is a suitable scalar multiple of  $\bar{\phi}_\tau^0(\tau \cdot)$ . Similarly an orthonormal basis of  $\bar{V}_\tau^1$  consists of a single function  $\bar{\phi}_\tau^1$  which is a suitable scalar multiple of  $\bar{\phi}_{\tau^2}^0(\tau \cdot)$ , and an orthonormal basis of  $\bar{V}_{\tau^2}^1$  consists of a single function  $\bar{\phi}_{\tau^2}^1$  which is a suitable scalar multiple of  $\bar{\phi}_{\tau^3}^0(\tau \cdot)$ .

For  $a = 1, \tau, \tau^2$ , let

$$\hat{w}_a := c_{\hat{w}_a} (\bar{\phi}_a^1 - \langle \bar{\phi}_a^1, \bar{\phi}_a^0 \rangle \bar{\phi}_a^0),$$

where  $c_{\hat{w}_a}$  is a constant so the  $\|\hat{w}_a\| = 1$ . By construction for each  $a > 0$

$$(54) \quad \hat{w}_a = \hat{w}_{\beta(a)}(\cdot - \mu(a))$$

forms an orthonormal basis for the one dimensional space  $\hat{W}_a$ .

In Figure 3 we see  $\hat{w}_1, \hat{w}_\tau$ , and  $\hat{w}_{\tau^2}$ .




FIGURE 3.  $\hat{w}_a$  for  $a = 1, \tau, \tau^2$

We now need to construct a basis for  $\tilde{W}_{\tau^2}$ . We begin by constructing the spaces defined in Section 2.1. A basis for  $T_{\tau^2}$  consists of the single function

$$t_{\tau^2} := \bar{\phi}_{\tau^2}^0 - \langle \bar{\phi}_{\tau^2}^0, \bar{\phi}_{\tau^2}^1 \rangle \bar{\phi}_{\tau^2}^1.$$

Since  $m_{\tau^2} = m_{\tau^2}^+ = m_{\tau^2}^- = 0$ , by equations (52) and (53), it follows from Lemma 13 that  $U_{\tau^2} = T_{\tau^2}$ . Thus  $S_{\tau^2}$  is spanned by the single function

$$s_{\tau^2} := (\chi_{[\tau^2, 1+\tau^2]} - \chi_{[\tau, \tau^2]})t_{\tau^2}.$$

By Lemma 16 an orthonormal basis of  $\tilde{W}_{\tau^2}$  consists of the single function

$$\tilde{w}_{\tau^2} := c_{\tilde{w}_{\tau^2}}(s_{\tau^2} - \langle s_{\tau^2}, \bar{\phi}_{\tau^2}^0 \rangle \bar{\phi}_{\tau^2}^0 - \langle s_{\tau^2}, \hat{w}_{\tau^2} \rangle \hat{w}_{\tau^2}),$$

where  $c_{\tilde{w}_{\tau^2}}$  is chosen so that  $\tilde{w}_{\tau^2}$  is of norm one. It follows that for  $b \in \mathbf{Z}_\tau^+$ , an orthonormal basis for  $\tilde{W}_{\tau^2+b\tau^3}$  consists of the single function

$$(55) \quad \tilde{w}_{\tau^2+b\tau^3} = \tilde{w}_{\tau^2}(\cdot - b\tau^3).$$

In Figure 4 we see  $\tilde{w}_{\tau^2}$ .




FIGURE 4.  $\tilde{w}_{\tau^2}$

To complete the wavelet construction, we must determine  $\check{W}_a$  for  $a \geq 0$ . Observe that

$$\check{k}_a^0 = \begin{cases} 3 & \text{if } a = 0 \\ 2 & \text{if } a > 0 \end{cases}$$

and

$$\check{k}_a^1 = \begin{cases} 6 & \text{if } a = 0 \\ 2 & \text{if } a \in 1 + \tau^2 \mathbf{Z}_\tau^+ \\ 5 & \text{if } a > 0 \in \tau \mathbf{Z}_\tau^+. \end{cases}$$

It follows from Theorem 11 and equations (52) and (53) that

$$\dim \check{W}_a = \begin{cases} 2 & \text{if } a = 0 \\ 0 & \text{if } a \in 1 + \tau^2 \mathbf{Z}_\tau^+ \\ 1 & \text{if } a > 0 \in \tau \mathbf{Z}_\tau^+. \end{cases}$$

We will be using equation (35) to construct the spaces  $\check{W}_a$ . We first remark that for any  $a > 0$ ,  $A_a^-$  is spanned by the single function

$$f_a^- := c_{f_a^-}(\chi_{[a-, a]}(\bar{\phi}_a^0 - \langle \bar{\phi}_a^0, \bar{\phi}_a^1 \rangle \bar{\phi}_a^1)),$$

and  $A_a^+$  is spanned by the single function

$$f_a^+ := c_{f_a^+}(\chi_{[a, a+]}(\bar{\phi}_a^0 - \langle \bar{\phi}_a^0, \bar{\phi}_a^1 \rangle \bar{\phi}_a^1)).$$

Here the constants  $c_{f_a^\pm}$  are chosen so that  $f_a^\pm$  are of norm one.

Since  $\dim \check{W}_\tau = 1$ , we need only construct one function in that space. We first observe that  $\bar{\phi}_{1+\tau^2}^0(\tau \cdot) \in \check{V}_\tau^1$ . Also,  $\check{V}_\tau^0$  is spanned by the functions  $q \circ \sigma_\tau$

and  $z^{1/\tau} \circ \sigma_\tau$ . Thus if we define  $\check{w}_\tau$  as the normalized form of the function  $(I - P_{\text{span}\{f_\tau^+, q \circ \sigma_\tau, z^{1/\tau} \circ \sigma_\tau, f_{\tau^2}^-\}}) \bar{\phi}_{1+\tau^2}^0(\tau \cdot)$ , then  $\check{w}_\tau$  forms an orthonormal basis for  $\check{W}_\tau$ .

It is easy to check that if we define  $\check{w}_{\tau^2} := \check{w}_\tau(\cdot - 1)$  then  $\check{w}_{\tau^2}$  forms an orthonormal basis for  $\check{W}_{\tau^2}$ .

Next, for  $b \in \mathbf{Z}_\tau^+$  we define

$$\begin{aligned}\check{w}_{\tau+\tau^2b} &:= \check{w}_\tau(\cdot - \tau^2b) \\ \check{w}_{\tau^2+\tau^3b} &:= \check{w}_{\tau^2}(\cdot - \tau^3b).\end{aligned}$$

It follows that for  $b \in \mathbf{Z}_\tau^+$ ,  $\check{w}_{\tau+\tau^2b}$  (resp.  $\check{w}_{\tau^2+\tau^3b}$ ) forms an orthonormal basis for  $\check{W}_{\tau+\tau^2b}$  (resp.  $\check{W}_{\tau^2+\tau^3b}$ ).

In Figure 5 we see  $\check{w}_\tau$ .




FIGURE 5.  $\check{w}_\tau$

It is similarly easy to check that  $\check{w}_{0,1} := \check{w}_\tau(\cdot + \tau) \in \check{W}_0$ . Since  $\dim \check{W}_0 = 2$  we need to construct one more function in this space. Using the the fact that  $\check{W}_0 = \check{V}_0^1 \ominus (\check{V}_0^0 \oplus A_1^-)$ , we can use the same type of argument as above to construct  $\check{w}_{0,2}$  so that  $\{\check{w}_{0,1}, \check{w}_{0,2}\}$  is an orthonormal basis for  $\check{W}_0$ .

In Figure 6 we see  $\check{w}_{0,2}$ .




FIGURE 6.  $\check{w}_{0,2}$

We now have that

$$\Psi_0 = \begin{bmatrix} \check{w}_{0,1} \\ \check{w}_{0,2} \end{bmatrix}, \Psi_1 = [\hat{w}_1], \Psi_\tau = \begin{bmatrix} \hat{w}_\tau \\ \check{w}_\tau \end{bmatrix}, \text{ and } \Psi_{\tau^2} = \begin{bmatrix} \hat{w}_{\tau^2} \\ \check{w}_{\tau^2} \end{bmatrix}.$$

For  $a > \tau^2$ ,  $\Psi_a = \Psi_{\beta(a)}(\cdot - \mu(a))$ .

Since  $S(\Phi^k) \subset S(\Phi^{k+1})$ , for  $k \in \mathbf{Z}$ , it follows from Theorem 7 that there exists an orthonormal basis,  $\Psi^k$  centered on  $\mathbf{a}^k$  so that  $S(\Phi^{k+1}) = S(\Phi^k) \oplus S(\Psi^k)$ .

For  $k \in \mathbf{Z}$  and  $a \in \mathbf{Z}_\tau^+$ ,

$$\Psi_{k,a} := \tau^{k/2} \Psi_a(\tau^k \cdot).$$

We now have that, for  $k \in \mathbf{Z}$  and  $b \in \mathbf{a}^k$ ,

$$\Psi_{b,\mathbf{a}^k}^k = \Psi_{k,\tau^k b}.$$

The wavelet coefficient matrices from section 2.3 can now be computed. For  $a \in \mathbf{Z}_\tau^+$  and  $a' \in \{a_-, a\}$ ,  $d_{aa'} = \langle \Psi_a, \Phi_{a'}^1 \rangle$ . In particular we have

$$\Psi_0 = d_{00} \Phi_0^1, \text{ and } \Psi_a = d_{aa_-} \Phi_{a_-}^1 + d_{aa} \Phi_a^1, \text{ for } a > 0.$$

Recall that

$$\Phi_0^1 = \begin{bmatrix} \Phi_{1,0} \\ \Phi_{1,1} \end{bmatrix}.$$

Thus

$$d_{00} = \langle \Psi_0, \Phi_0^1 \rangle = [\langle \Psi_0, \Phi_{1,0} \rangle \quad \langle \Psi_0, \Phi_{1,1} \rangle].$$

This suggests defining, for  $a, a' \in \mathbf{Z}_\tau^+$ ,

$$D_{a,a'} := \langle \Psi_a, \Phi_{1,a'} \rangle.$$

We now have

$$d_{00} = \begin{bmatrix} D_{0,0} & D_{0,1} \end{bmatrix} \text{ and } d_{10} = \begin{bmatrix} D_{1,0} & D_{1,1} \end{bmatrix}.$$

Similarly it follows that  $d_{11} = D_{1,\tau}$ ,  $d_{\tau 1} = 0$ ,

$$d_{\tau\tau} = \begin{bmatrix} D_{\tau,\tau^2} & D_{\tau,1+\tau^2} \end{bmatrix}, d_{\tau^2\tau} = \begin{bmatrix} D_{\tau^2,\tau^2} & D_{\tau^2,1+\tau^2} \end{bmatrix},$$

and

$$d_{\tau^2\tau^2} = \begin{bmatrix} D_{\tau^2,\tau^3} & D_{\tau^2,1+\tau^3} \end{bmatrix}.$$

For  $a > \tau^2$ ,  $d_{aa} = d_{\beta(a)\beta(a)}$  and  $d_{aa_-} = d_{\beta(a)\beta(a)_-}$ .

In Table 2 we provide the matrices  $D_{a,a'}$ , mentioned above, which were computed with the aid of *Mathematica*<sup>TM</sup>.

TABLE 2. Wavelet Coefficients for  $\Psi$

$$\begin{aligned} D_{0,0} &= \begin{pmatrix} 0 & \frac{1}{2\sqrt{300+134\sqrt{5}}} & -\frac{1}{2}\sqrt{\frac{1}{10}(-2+\sqrt{5})} \\ \frac{1}{2}(-1+\sqrt{5}) & \frac{1}{2\sqrt{2(5096+2279\sqrt{5})}} & -\frac{1}{2}\sqrt{-168+\frac{761}{2\sqrt{5}}} \end{pmatrix} \\ D_{0,1} &= \begin{pmatrix} \frac{1}{55}(-5+18\sqrt{5}) & -\frac{1}{2}\sqrt{-\frac{1}{2}+\frac{2}{\sqrt{5}}} & -\frac{6+7\sqrt{5}}{22\sqrt{2}} \\ -\frac{1}{11}\sqrt{\frac{1}{10}(6085-2689\sqrt{5})} & -\frac{1}{2}\sqrt{\frac{11}{2(237+106\sqrt{5})}} & -\frac{19}{2}\sqrt{\frac{5}{2(7925+3538\sqrt{5})}} \end{pmatrix} \\ D_{1,0} &= \begin{pmatrix} 0 & \frac{1}{\sqrt{445+199\sqrt{5}}} & -\sqrt{\frac{11}{5(69+31\sqrt{5})}} \end{pmatrix} \\ D_{1,1} &= \begin{pmatrix} \sqrt{\frac{2}{55}(13-5\sqrt{5})} & -\frac{1}{2}\sqrt{-45+\frac{103}{\sqrt{5}}} & \frac{1}{2}\sqrt{\frac{1}{11}(263-113\sqrt{5})} \end{pmatrix} \\ D_{1,\tau} &= \begin{pmatrix} \sqrt{-13+6\sqrt{5}} & 0 & 0 \end{pmatrix} \\ D_{\tau,\tau^2} &= \begin{pmatrix} \sqrt{-\frac{5}{8}+\frac{3\sqrt{5}}{8}} & \frac{1}{4}\sqrt{\frac{31}{62071+27759\sqrt{5}}} & -\frac{1}{8}\sqrt{\frac{4141-\frac{9161}{\sqrt{5}}}{5}} \\ 0 & \frac{1}{2\sqrt{300+134\sqrt{5}}} & -\frac{1}{2}\sqrt{\frac{1}{10}(-2+\sqrt{5})} \end{pmatrix} \end{aligned}$$

Table 2 continued

$$\begin{aligned}
D_{\tau, \tau^2+1} &= \begin{pmatrix} -\sqrt{-\frac{7331}{968} + \frac{16557}{968\sqrt{5}}} & -\frac{1}{4}\sqrt{\frac{341}{2887+1291\sqrt{5}}} & -\frac{19}{4}\sqrt{\frac{155}{96375+43163\sqrt{5}}} \\ \frac{1}{55}(-5+18\sqrt{5}) & -\frac{1}{2}\sqrt{-\frac{1}{2} + \frac{2}{\sqrt{5}}} & -\frac{6+7\sqrt{5}}{22\sqrt{2}} \end{pmatrix} \\
D_{\tau^2, \tau^2} &= \begin{pmatrix} 0 & \sqrt{\frac{873}{62} - \frac{976}{31\sqrt{5}}} & -\frac{11}{\sqrt{6770+3040\sqrt{5}}} \\ 0 & \frac{1}{2\sqrt{199+89\sqrt{5}}} & -\frac{1}{4}\sqrt{31 - \frac{69}{\sqrt{5}}} \\ 0 & 0 & 0 \end{pmatrix} \\
D_{\tau^2, \tau^2+1} &= \begin{pmatrix} \sqrt{\frac{2}{155}(23-9\sqrt{5})} & -\sqrt{-\frac{221}{31} + \frac{1009}{62\sqrt{5}}} & \sqrt{\frac{1}{62}(234-101\sqrt{5})} \\ \sqrt{-\frac{5}{22} + \frac{13}{22\sqrt{5}}} & \frac{9-5\sqrt{5}}{4\sqrt{2}} & \sqrt{-\frac{565}{176} + \frac{263\sqrt{5}}{176}} \\ 0 & 0 & 0 \end{pmatrix} \\
D_{\tau^2, \tau^3} &= \begin{pmatrix} \sqrt{\frac{1}{31}(7+4\sqrt{5})} & \sqrt{\frac{5}{2(397555+177792\sqrt{5})}} & -\frac{121}{\sqrt{52754+23600\sqrt{5}}} \\ 0 & -\frac{1}{2\sqrt{9349+4181\sqrt{5}}} & \frac{121}{2}\sqrt{\frac{11}{34145+15249\sqrt{5}}} \\ 0 & \frac{1}{2\sqrt{300+134\sqrt{5}}} & -\frac{1}{2}\sqrt{\frac{1}{10}(-2+\sqrt{5})} \end{pmatrix} \\
D_{\tau^2, \tau^3+1} &= \begin{pmatrix} -38\sqrt{\frac{2}{94393+42201\sqrt{5}}} & -\sqrt{\frac{55}{2(18490+8269\sqrt{5})}} & -19\sqrt{\frac{5}{2(123546+55249\sqrt{5})}} \\ 19\sqrt{\frac{2}{5545+2483\sqrt{5}}} & \frac{11}{2\sqrt{4783+2139\sqrt{5}}} & \frac{19}{2}\sqrt{\frac{5}{14525+6497\sqrt{5}}} \\ \frac{1}{55}(-5+18\sqrt{5}) & -\frac{1}{2}\sqrt{-\frac{1}{2} + \frac{2}{\sqrt{5}}} & -\frac{6+7\sqrt{5}}{22\sqrt{2}} \end{pmatrix}
\end{aligned}$$

## REFERENCES

- [1] D. Bruff, Wavelets on nonuniform knot sequences, Vanderbilt University Ph.D. Thesis, (2003).
- [2] D. Bruff and D. P. Hardin, Squeezable bases and semi-regular multiresolutions, *Wavelet analysis* (Hong Kong, 2001), (2002), 9–22.
- [3] M. Charina and J. Stöckler, Tight wavelet frames for irregular multiresolution analysis, *ACHA*, **25**, (2008), 89–113.
- [4] C.K. Chui and X. Shi, Orthonormal wavelets and tight frames with arbitrary real dilations, *Appl. Comput. Harmon. Anal.* **9**, (2000), 243–264.
- [5] A. Cohen, and N. Dyn, Nonstationary subdivision schemes, multiresolution analysis, and wavelet packets, *Wavelet Anal. Appl.*, (1998), **7**, 189–200.
- [6] I. Daubechies, Orthonormal bases of compactly supported wavelets, *Comm. Pure Appl. Math.*, **41** (1988), 909–996.
- [7] I. Daubechies, I. Guskov, P. Schröder, and W. Sweldens, Wavelets on Irregular Point Sets, *Phil. Trans. R. Soc. Lond. A*, **357**, (1999), 2397–2413.
- [8] I. Daubechies, I. Guskov, and W. Sweldens, Regularity of Irregular Subdivision, *Const. approx.*, **15**, (1999), 381–426.
- [9] G. C. Donovan, J. S. Geronimo, and D. P. Hardin, Intertwining multiresolution analyses and the construction of piecewise polynomial wavelets, *SIAM J. Math. Analysis* **27**, (1996), 1791–1815.
- [10] G. C. Donovan, J. S. Geronimo, and D. P. Hardin, Squeezable orthogonal bases and adaptive least squares, *Wavelet Applications in Signal and Image Processing V*, Aldroubi, Laine & Unser, editors, SPIE Conf. Proc. **3169**, (1997), 48–54.
- [11] G. Donovan, J. Geronimo, and D. Hardin, Squeezable Orthogonal Bases: Accuracy and Smoothness, *SIAM J. Numer. Anal.*, **40**, (2002), 1077–1099.
- [12] G. Donovan, J. Geronimo, and D. Hardin, Orthogonal polynomials and the construction of piecewise polynomial smooth wavelets, *SIAM J. Math. Anal.* **30** (1999), 1029–1056.
- [13] J. S. Geronimo, D. P. Hardin, and P. R. Massopust, Fractal functions and wavelet expansions based on several scaling functions, *J. Approx. Theory* **78**, (1994), 373–401.
- [14] J-P Gazeau and J. Patera, Tau-wavelets of Haar, *J. Phys. A: Math. Gen.*, **29**, (1996), 4549–4559.
- [15] J-P Gazeau and V. Spiridonov, Toward discrete wavelets with irrational scaling factor, *J. Math. Phys.*, **37**, (1996), 3001–3013.

- [16] C. Herley, J. Kovačević, and M. Vetterli, Tilings of the time-frequency plane: Construction of arbitrary orthogonal bases and fast tiling algorithms. *IEEE Trans Sig. Proc*, **41**, (1993), 3341-3359
- [17] E. Hernandez, X. Wang, and G. Weiss, Smoothing minimally supported frequency (MSF) wavelets: Part I *J. Fourier Anal. Appl*, **2**, (1995), 239-340.
- [18] E. Hernandez, X. Wang, and G. Weiss, Smoothing minimally supported frequency wavelets: Part II *J. Fourier Anal. Appl*, **3**, (1997), 23-41.
- [19] M. Lothaire, Algebraic Combinatorics on Words, Cambridge University Press, Cambridge, 2002.
- [20] L. Schumaker, Spline Functions: Basic Theory, third edition, Cambridge University Press, Cambridge, 2007.
- [21] W. Sweldens, The lifting scheme: a construction of second generation wavelets, *SIAM J. Math. Anal*, **29**, (1997), 511-546.